

## APPLICATION OF THE MEAN ERGODIC THEOREM TO CERTAIN SEMIGROUPS

Gerd Herzog and Christoph Schmoeger

*Communicated by*

ABSTRACT. We study the asymptotic behaviour of solutions of the Cauchy problem  $u' = (\sum_{j=1}^n (A_j + A_j^{-1}) - 2nI)u$ ,  $u(0) = x$  as  $t \rightarrow \infty$ , for invertible isometries  $A_1, \dots, A_n$ .

### 1. Introduction

Let  $E$  be a complex Banach space,  $L(E)$  the Banach algebra of all bounded linear operators on  $E$ , and let  $A_1, \dots, A_n \in L(E)$  be invertible, pairwise commuting, and such that  $\|A_k\| = \|A_k^{-1}\| = 1$  ( $k = 1, \dots, n$ ). Let  $T_1, \dots, T_n \in L(E)$  be defined by  $T_k = A_k + A_k^{-1} - 2I$ , and let  $T = T_1 + \dots + T_n$ . The aim of this paper is to clear the asymptotic behaviour of the Cauchy problem

$$(1.1) \quad u'(t) = Tu(t), \quad u(0) = u_0$$

that is of  $t \mapsto \exp(tT)u_0$  for  $t \rightarrow \infty$ . Such problems occur in a natural way by semidiscretization of the parabolic Cauchy problem  $v_t = \Delta v$ ,  $v(0, x) = v_0(x)$ : For example, if  $v_0 : \mathbb{R}^n \rightarrow \mathbb{R}$  is bounded, the longitudinal line method, see for example [4], with step size 1 leads to a linear Cauchy problem of type (1.1) in  $l^\infty(\mathbb{Z}^n)$  with

$$A_k x = (x(j_1, j_2, \dots, j_{k-1}, j_k + 1, j_{k+1}, \dots, j_n))_{j \in \mathbb{Z}^n}.$$

The corresponding problem for the heat equation was studied in [1].

### 2. Notations and preliminaries

For  $A \in L(E)$  let  $N(A)$ ,  $A(E)$ ,  $\sigma(A)$  and  $r(A)$  denote the kernel, the range, the spectrum and the spectral radius of  $A$ , respectively. Let  $\mathbb{D}$  denote the complex unit circle  $\{z \in \mathbb{C} : |z| < 1\}$ .

PROPOSITION 2.1. *Let  $A \in L(E)$ ,  $0 \notin \sigma(A)$  and  $\|A\| = \|A^{-1}\| = 1$ . Then:*

- (1)  *$A$  is an isometry;*

- (2)  $\|A^n\| = \|A\|^n = 1$  ( $n \in \mathbb{N}$ ), hence  $A$  is normaloid;
- (3)  $r(A) = 1$  and  $\overline{\sigma(A)} \subseteq \partial\mathbb{D}$ ;
- (4)  $N(A - I) \cap \overline{(A - I)(E)} = \{0\}$ ;
- (5)  $(A - I)(E) = (A^{-1} - I)(E)$ ;
- (6)  $N(A - I) = N((A - I)^2)$ ;
- (7)  $N(A - I) \oplus \overline{(A - I)(E)}$  is closed;
- (8) if  $(A - I)(E)$  is closed then  $E = N(A - I) \oplus \overline{(A - I)(E)}$ .

PROOF. (1) and (2) are obvious.

(3): From (2) we get  $r(A) = 1$ . Next, it is clear that

$$\sigma(A) \cup \sigma(A^{-1}) \subseteq \overline{\mathbb{D}}.$$

Since  $\sigma(A) = \{z \in \mathbb{C} : z^{-1} \in \sigma(A^{-1})\}$ , we conclude  $\sigma(A) \subseteq \partial\mathbb{D}$ .

(4): Let  $x \in N(A - I) \cap \overline{(A - I)(E)}$ , let  $\varepsilon > 0$  and choose  $z \in E$  such that  $\|x - (A - I)z\| < \varepsilon$ . According to [2, Satz 102.3], we have  $\|x\| \leq \|x - (A - I)z\| < \varepsilon$ , hence  $x = 0$ .

(5): Follows from  $(A - I)x = (I - A^{-1})(Ax)$ .

(6): Follows from [2, Satz 102.3].

(7): Choose a sequence  $(x_n)$  in  $N(A - I) \oplus \overline{(A - I)(E)}$  with  $x_n \rightarrow x_0$  and corresponding decompositions  $x_n = y_n + z_n$ . According to [2, Satz 102.3] we have

$$\|y_n - y_m\| \leq \|x_n - x_m\| \quad (n, m \in \mathbb{N}),$$

hence  $(y_n)$  is convergent to a vector  $y_0 \in N(A - I)$ . Thus

$$z_n = x_n - y_n \rightarrow x_0 - y_0 \in \overline{(A - I)(E)},$$

and therefore  $x_0 \in N(A - I) \oplus \overline{(A - I)(E)}$ .

(8): Follows from [2, Satz 72.4 and 102.4].  $\square$

PROPOSITION 2.2. Let  $A \in L(E)$  be as in Proposition 2.1, let  $T = A + A^{-1} - 2I$ , and let  $c : [0, \infty) \rightarrow \mathbb{R}$  denote the function

$$c(t) = \exp(-t) \left( 1 + \sum_{n=0}^{\infty} \frac{t^n}{n!} \left| 1 - \frac{t}{n+1} \right| \right).$$

We have

- (1)  $\|\exp(tT)\| \leq 1$  ( $t \geq 0$ );
- (2)  $t \mapsto \sqrt{t}c(t)$  is bounded on  $[0, \infty)$  and

$$\|\exp(tT)(A - I)x\| \leq c(t)\|x\| \quad (t \geq 0, x \in E);$$

- (3)  $\lim_{t \rightarrow \infty} \exp(tT)y = 0$  ( $y \in (A - I)(E)$ );
- (4) if  $y \in E$  then

$$\lim_{t \rightarrow \infty} \exp(tT)y = 0 \iff y \in \overline{(A - I)(E)};$$

- (5)  $N(A - I) = \{x \in E : \exp(tT)x = x \ (t \geq 0)\}$ .

PROOF. (1): For each  $t \geq 0$

$$\begin{aligned} \|\exp(tT)\| &= \|\exp(-2t) \exp(tA) \exp(tA^{-1})\| \\ &\leq \exp(-2t) \exp(t\|A\|) \exp(t\|A^{-1}\|) = 1. \end{aligned}$$

(2): Since  $T = (A^{-1} - I) + (A - I)$  we have

$$\exp(tT)(A - I) = \exp(t(A^{-1} - I)) \exp(t(A - I))(A - I) \quad (t \in \mathbb{R}),$$

and

$$\begin{aligned} \exp(t(A - I))(A - I)x &= \exp(-t) \sum_{n=0}^{\infty} \frac{t^n}{n!} (A^{n+1} - A^n)x \\ &= \exp(-t) \left( \sum_{n=0}^{\infty} \frac{t^n}{n!} \left(1 - \frac{t}{n+1}\right) \right) A^{n+1}x - \exp(-t)x. \end{aligned}$$

Hence, since  $\|A\| = 1$ ,

$$\begin{aligned} \|\exp(tT)(A - I)x\| &\leq \|\exp(t(A^{-1} - I))\| \|\exp(t(A - I))(A - I)x\| \\ &\leq c(t)\|x\| \quad (t \geq 0, x \in E). \end{aligned}$$

To see that  $t \mapsto \sqrt{t}c(t)$  is bounded on  $[0, \infty)$  let  $N \in \mathbb{N}$  and  $N \leq t \leq N + 1$ . Then

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{t^n}{n!} \left|1 - \frac{t}{n+1}\right| &= \sum_{n=0}^{N-1} \frac{t^n}{n!} \left(\frac{t}{n+1} - 1\right) + \sum_{n=N}^{\infty} \frac{t^n}{n!} \left(1 - \frac{t}{n+1}\right) \\ &= 2\frac{t^N}{N!} - 1, \end{aligned}$$

and therefore

$$\begin{aligned} \sqrt{t}c(t) &= \sqrt{t} \exp(-t) \left(1 + \sum_{n=0}^{\infty} \frac{t^n}{n!} \left|1 - \frac{t}{n+1}\right|\right) \\ &\leq \sqrt{N+1} \exp(-N) \left(1 + 2\frac{(N+1)^N}{N!} - 1\right) \\ &= \frac{2(N+1)^{N+1/2} \exp(-N)}{N!}, \end{aligned}$$

which is bounded according to Stirling's formula.

(3): Follows from (2).

(4): The implication  $\Leftarrow$  follows from (3). Now suppose that  $\exp(tT)y \rightarrow 0$  as  $t \rightarrow \infty$ . Since

$$\begin{aligned} \exp(tT)y &= y + \sum_{n=1}^{\infty} \frac{t^n}{n!} T^n y \\ &= y + (A - I) \sum_{n=1}^{\infty} \frac{t^n}{n!} (A - I)^{n-1} (I - A^{-1})^n y \end{aligned}$$

we conclude  $y \in \overline{(A - I)(E)}$ .

(5): The inclusion

$$N(A - I) \subseteq \{x \in E : \exp(tT)x = x \ (t \geq 0)\}$$

is obvious. Now suppose that  $x \in E$  and  $\exp(tT)x = x$  ( $t \geq 0$ ). By differentiation  $0 = T \exp(tT)x$  ( $t \geq 0$ ), thus  $A^{-1}(A - I)^2x = Tx = 0$ . Part (6) of Proposition 2.1 now shows that  $x \in N(A - I)$ .  $\square$

### 3. The asymptotic behaviour of $\exp(tT)$

**THEOREM 3.1.** *Let  $A$  and  $T$  be as in Proposition 2.1. For  $x \in E$  the following assertions are equivalent:*

- (1)  $\lim_{t \rightarrow \infty} \exp(tT)x$  exists in  $E$  [resp.  $\lim_{t \rightarrow \infty} \exp(tT)x = 0$ ];
- (2)  $x \in N(A - I) \oplus \overline{(A - I)(E)}$  [resp.  $x \in \overline{(A - I)(E)}$ ];
- (3) the sequence

$$\left( \frac{x + Ax + \cdots + A^m x}{m + 1} \right)_{m \in \mathbb{N}}$$

is convergent [resp. is convergent with limit 0].

**PROOF.** That (2) implies (1) follows from Proposition 2.2. Now, assume that (1) holds, and let  $z = \lim_{t \rightarrow \infty} \exp(tT)x$ . As in the proof of part (4) of Proposition 2.2

$$\exp(tT)x = x + (A - I) \sum_{n=1}^{\infty} \frac{t^n}{n!} (A - I)^{n-1} (I - A^{-1})^n x,$$

hence  $x - z \in \overline{(A - I)(E)}$ . [In particular, if  $z = 0$  then  $x \in \overline{(A - I)(E)}$ .] From part (3) of Proposition 2.2 we obtain

$$(A - I)z = \lim_{t \rightarrow \infty} \exp(tT)(A - I)x = 0,$$

and therefore  $x = z + (x - z) \in N(A - I) \oplus \overline{(A - I)(E)}$ .

The equivalence of (2) and (3) is proved in [3, Ch.2, Theorem 1.3].  $\square$

According to part (8) of Proposition 2.1 the following corollary shows, that  $\lim_{t \rightarrow \infty} \exp(tT)x$  exists for each  $x \in E$  if  $T(E)$  is closed:

**COROLLARY 3.1.** *We have*

- (1)  $T(E) = (A - I)^2(E) \subseteq (A - I)(E) \subseteq \overline{T(E)}$ ;
- (2)  $\frac{T(E)}{(A - I)(E)} = \frac{\overline{T(E)}}{\overline{(A - I)(E)}} \iff (A - I)^2(E) = (A - I)(E) \iff (A - I)(E) = \overline{(A - I)(E)}$ .

**PROOF.** (1): Part (5) of Proposition 2.1 gives

$$T(E) = (A - I)^2(E) \subseteq (A - I)(E).$$

As in the proof of Theorem 3.1 we obtain  $(A - I)(E) \subseteq \overline{T(E)}$ . Now, (2) follows by [2, Satz 102.4].  $\square$

#### 4. The general case

Now, let  $A_1, \dots, A_n, T_1, \dots, T_n$  and  $T$  be as in section 1. Moreover we introduce the following subspaces of  $E$ :

$$X_1 = \bigcap_{j=1}^n N(A_j - I), \quad X_2 = \overline{\sum_{j=1}^n (A_j - I)(E)}, \quad X = X_1 + X_2.$$

THEOREM 4.1. *Under the assumptions above*

- (1)  $X_2 = \{x \in E : \lim_{t \rightarrow \infty} \exp(tT)x = 0\}$ ;
- (2)  $X_1 = \{x \in E : \exp(tT)x = x \ (t \geq 0)\}$ ;
- (3)  $X = \{x \in E : \lim_{t \rightarrow \infty} \exp(tT)x \text{ exists in } E\}$ ;
- (4)  $X_1 \cap X_2 = \{0\}$ , and  $X$  is closed.

PROOF. (1): Let  $x \in X_2$ . Then  $x = \lim_{m \rightarrow \infty} x_m$  where

$$x_m \in \sum_{j=1}^n (A_j - I)(E).$$

By part (1) and part (3) of Proposition 2.2 we obtain

$$\lim_{t \rightarrow \infty} \exp(tT)x_m = 0 \quad (m \in \mathbb{N}).$$

Let  $\varepsilon > 0$ , and choose  $N \in \mathbb{N}$  such that  $\|x - x_N\| < \varepsilon/2$ . Next, choose  $t_0 \in [0, \infty)$  such that  $\|\exp(tT)x_N\| < \varepsilon/2$  ( $t \geq t_0$ ). Then

$$\begin{aligned} \|\exp(tT)x\| &= \|\exp(tT)(x - x_N) + \exp(tT)x_N\| \\ &\leq \|x - x_N\| + \|\exp(tT)x_N\| < \varepsilon \quad (t \geq t_0). \end{aligned}$$

Thus  $\lim_{t \rightarrow \infty} \exp(tT)x = 0$ .

Now suppose that  $x \in E$  and  $\lim_{t \rightarrow \infty} \exp(tT)x = 0$ . Set

$$h(t) = \sum_{n=1}^{\infty} \frac{t^n}{n!} T^n x.$$

Since  $T_j = (A_j - I)(I - A_j^{-1})$  ( $j = 1, \dots, n$ ), we have

$$Tx = \sum_{j=1}^n (A_j - I)(I - A_j^{-1})x \in \sum_{j=1}^n (A_j - I)(E),$$

hence

$$h(t) \in \sum_{j=1}^n (A_j - I)(E).$$

Thus,  $\exp(tT)x = x + h(t)$  and  $\lim_{t \rightarrow \infty} \exp(tT)x = 0$  imply  $x \in X_2$ .

(2): The inclusion  $\subseteq$  is obvious. For the reversed inclusion let  $x \in E$  be such that  $\exp(tT)x = x$  ( $t \geq 0$ ). Then by part (1) we obtain

$$(A_j - I)x = \exp(tT)(A_j - I)x \rightarrow 0 \quad (t \rightarrow \infty) \quad (j = 1, \dots, n),$$

hence  $x \in X_1$ .

(3): Here, the inclusion  $\subseteq$  follows from parts (1) and (2) directly. Now, assume that  $x \in E$  is such that  $\lim_{t \rightarrow \infty} \exp(tT)x = z$ . As in the proof of part (1)

$$\exp(tT)x = x + h(t), \quad h(t) \in X_2.$$

Therefore  $x - z \in X_2$ . From part (1) we derive

$$(A_j - I)z = \lim_{t \rightarrow \infty} \exp(tT)(A_j - I)x = 0 \quad (j = 1, \dots, n).$$

Thus  $z \in X_1$ , and so  $x = z + (x - z) \in X_1 \oplus X_2 = X$ .

(4): Let  $x \in X_1 \cap X_2$ . Then, by parts (1) and (2), we have

$$\exp(tT)x = x \quad (t \geq 0), \quad \exp(tT)x \rightarrow 0 \quad (t \rightarrow \infty),$$

thus  $x = 0$ . Next, if  $(x_m)$  is a sequence in  $X$  with limit  $x_0$ , then there exist sequences  $(y_m)$  and  $(z_m)$  in  $X_1$  and  $X_2$ , respectively, with  $x_m = y_m + z_m$ . From part (1) and part (2) we obtain

$$\exp(tT)(x_m - x_k) \rightarrow y_m - y_k \quad (t \rightarrow \infty).$$

Since  $\|\exp(tT)(x_m - x_k)\| \leq \|x_m - x_k\|$  ( $t \geq 0$ ), we have

$$\|y_m - y_k\| \leq \|x_m - x_k\|,$$

thus  $(y_m)$  is convergent. Let  $y_0 = \lim_{m \rightarrow \infty} y_m$ . Then  $z_m = x_m - y_m \rightarrow x_0 - y_0$ . Hence we have  $y_0 \in X_1$ ,  $x_0 - y_0 \in X_2$ , and therefore  $x_0 \in X_1 \oplus X_2 = X$ .  $\square$

The following result provides sufficient conditions for the convergence of  $\exp(tT)x$ .

**THEOREM 4.2.** *Let  $(k_1, \dots, k_n) \in \mathbb{N}_0^n$ , and set  $B = A_1^{k_1} \dots A_n^{k_n}$ . We have*

- (1)  $\bigcap_{j=1}^n \left( N(A_j - I) \oplus \overline{(A_j - I)(E)} \right) \subseteq X$ ;
- (2)  $\overline{(B - I)(E)} \subseteq X_2$ ;
- (3) if  $x \in E$  and if the sequences

$$\left( \frac{x + A_j x + \dots + A_j^m x}{m + 1} \right)_{m \in \mathbb{N}}$$

are convergent ( $j = 1, \dots, n$ ), then  $\lim_{t \rightarrow \infty} \exp(tT)x$  exists in  $E$ ;

- (4) if  $x \in E$  and if the sequence

$$\left( \frac{x + Bx + \dots + B^m x}{m + 1} \right)_{m \in \mathbb{N}}$$

is convergent to 0 in  $E$ , then  $\lim_{t \rightarrow \infty} \exp(tT)x = 0$ .

**PROOF.** According to Theorem 3.1 we see that (3) follows from (1), and (4) follows from (2).

For the proof of (1) we use induction. If  $n = 1$  the result follows by Theorem 3.1.

Suppose that  $n \in \mathbb{N}$  and that (1) holds. In the case of  $n + 1$  operators  $T_1, \dots, T_{n+1}$  we write  $T_0 = T_1 + \dots + T_n$ , so  $T = T_0 + T_{n+1}$ . Let

$$x \in \bigcap_{j=1}^{n+1} \left( N(A_j - I) \oplus \overline{(A_j - I)(E)} \right).$$

Then

$$x \in \bigcap_{j=1}^n \left( N(A_j - I) \oplus \overline{(A_j - I)(E)} \right), \quad x \in N(A_{n+1} - I) \oplus \overline{(A_{n+1} - I)(E)},$$

and therefore the limits  $\lim_{t \rightarrow \infty} \exp(tT_0)x$  and  $\lim_{t \rightarrow \infty} \exp(tT_{n+1})x$  exist in  $E$ . From

$$\begin{aligned} & \| \exp(tT)x - \exp(sT)x \| = \| \exp(tT_0) \exp(tT_{n+1})x - \exp(sT_0) \exp(sT_{n+1})x \| \\ & = \| \exp(tT_0)(\exp(tT_{n+1}) - \exp(sT_{n+1}))x + \exp(sT_{n+1})(\exp(tT_0) - \exp(sT_0))x \| \\ & \leq \| \exp(tT_{n+1})x - \exp(sT_{n+1})x \| + \| \exp(tT_0)x - \exp(sT_0)x \| \end{aligned}$$

we see that  $\lim_{t \rightarrow \infty} \exp(tT)x$  exists.

Next, we prove (2) for  $(k_1, \dots, k_n) \in \mathbb{N}^n$ , without loss of generality. Let  $p(z) = z_1^{k_1} \dots z_n^{k_n} - 1$  ( $z = (z_1, \dots, z_n)$ ), and note that there are polynomials  $q_1, \dots, q_n \in \mathbb{C}[z_1, \dots, z_n]$  such that

$$p(z) = (z_1 - 1)q_1(z) + \dots + (z_n - 1)q_n(z).$$

Hence

$$(B - I)x \in \sum_{j=1}^n (A_j - I)(E) \quad (x \in E),$$

and therefore  $\overline{(B - I)(E)} \subseteq X_2$ .  $\square$

## 5. Example

Let us return to the semidiscretization of  $v_t = \Delta v$  in  $\mathbb{R}^2$ , that is we consider  $E = l^\infty(\mathbb{Z}^2)$  and

$$A_1 x = (x(i+1, j))_{(i,j) \in \mathbb{Z}^2}, \quad A_2 x = (x(i, j+1))_{(i,j) \in \mathbb{Z}^2}.$$

Let  $k_1, k_2 \in \mathbb{N}$ , and assume that  $x \in l^\infty(\mathbb{Z}^2)$  is such that the sequence

$$\left( \left( \frac{x(i, j) + x(i+k_1, j+k_2) + \dots + x(i+mk_1, j+mk_2)}{m+1} \right)_{(i,j) \in \mathbb{Z}^2} \right)_{m \in \mathbb{N}}$$

tends to 0 as  $m \rightarrow \infty$  in  $l^\infty(\mathbb{Z}^2)$ . Then

$$\exp(tT)x \rightarrow 0 \quad (t \rightarrow \infty)$$

(apply part (4) of Theorem 4.2 with  $B = A_1^{k_1} A_2^{k_2}$ ).

### References

- [1] G. Herzog, R. Lemmert, *A note on discrete heat conduction*, Arch. Math. **75** (2000), 147-152..
- [2] H. Heuser, *Funktionalanalysis*, 2. Auflage. Mathematische Leitfäden. Stuttgart: Teubner (1986).
- [3] U. Krengel, *Ergodic theorems*. De Gruyter Studies in Mathematics, 6. Berlin-New York: Walter de Gruyter. VIII (1985).
- [4] W. Walter, *Some new aspects of the line method for parabolic differential equations*, Lect. Notes Math. **280** (1972), 181-189.

INSTITUT FÜR ANALYSIS, UNIVERSITÄT KARLSRUHE, 76128 KARLSRUHE, GERMANY  
*E-mail address:* `Gerd.Herzog@math.uni-karlsruhe.de`

INSTITUT FÜR ANALYSIS, UNIVERSITÄT KARLSRUHE, 76128 KARLSRUHE, GERMANY  
*E-mail address:* `christoph.schmoeger@math.uni-karlsruhe.de`