

Perturbation properties of some classes of operators

C. SCHMOEGER

RIASSUNTO: Sia X uno spazio di Banach complesso e $\mathcal{L}(X)$ l'algebra di Banach di tutti gli operatori lineari limitati in X . Considerate le seguenti famiglie di operatori:

$$\mathcal{D}(X) = \{T \in \mathcal{L}(X) : T(X) \text{ is closed and } N(T) \subseteq \bigcap_{n=1}^{\infty} T^n(X)\},$$

$$\mathcal{S}(X) = \{T \in \mathcal{D}(X) : T \text{ is relatively regular}\}.$$

si determinano i punti interni di $\mathcal{D}(X)$ e $\mathcal{S}(X)$, si dimostrano inoltre alcuni teoremi di perturbazione.

ABSTRACT: Let X be a complex Banach space and $\mathcal{L}(X)$ the Banach algebra of all bounded linear operators on X . We consider the following classes of operators:

$$\mathcal{D}(X) = \{T \in \mathcal{L}(X) : T(X) \text{ is closed and } N(T) \subseteq \bigcap_{n=1}^{\infty} T^n(X)\},$$

$$\mathcal{S}(X) = \{T \in \mathcal{D}(X) : T \text{ is relatively regular}\}.$$

We determine the interior points of $\mathcal{D}(X)$ and $\mathcal{S}(X)$ and prove some perturbation theorems.

1 – Introduction and terminology

Throughout this paper X denotes a Banach space over the complex

KEY WORDS AND PHRASES: *Relatively regular operators – Atkinson operators – Semi-Fredholm operators*

A.M.S. CLASSIFICATION: 47A53 – 47A99

field \mathbb{C} and $\mathcal{L}(X)$ the Banach algebra of all bounded linear operators on X . If $T \in \mathcal{L}(X)$ we denote by $N(T)$ the kernel of T and by $\alpha(T)$ the dimension of $N(T)$. The range of T is denoted by $T(X)$ and we define $\beta(T) = \text{codim} T(X)$.

$T \in \mathcal{L}(X)$ is called relatively regular if $TST = T$ for some $S \in \mathcal{L}(X)$. $\mathcal{R}(X)$ will denote the set of all relatively regular operators.

We shall make use of the following results [1, p. 10]:

1. $T \in \mathcal{R}(X)$ if and only if $N(T)$ and $T(X)$ are closed complemented subspaces of X .
2. If $TST = T$ for some $S \in \mathcal{L}(X)$, then TS is a projection onto $T(X)$ and $I - ST$ is a projection onto $N(T)$.

An operator T is called an Atkinson operator if $T \in \mathcal{R}(X)$ and at least one of $\alpha(T)$, $\beta(T)$ is finite. The set of Atkinson operators will be denoted by $\mathcal{A}(X)$.

We write $\mathcal{C}(X)$ for the set of operators having closed range. The class of semi-Fredholm operators is defined by

$$\mathcal{SF}(X) = \{T \in \mathcal{C}(X) : \alpha(T) < \infty \text{ or } \beta(T) < \infty\}.$$

We have $\mathcal{A}(X) \subseteq \mathcal{SF}(X)$. The index of $T \in \mathcal{SF}(X)$ is given by $\text{ind}(T) = \alpha(T) - \beta(T)$.

The following result is well known (for proofs see [1] and [3]).

THEOREM 1. *Let $T \in \mathcal{A}(X)$ (resp. $T \in \mathcal{SF}(X)$). Then there exists $\delta > 0$ such that*

- (a) $T - B \in \mathcal{A}(X)$ (resp. $T - B \in \mathcal{SF}(X)$), $\alpha(T - B) \leq \alpha(T)$, $\beta(T - B) \leq \beta(T)$ and $\text{ind}(T - B) = \text{ind}(T)$ for all $B \in \mathcal{L}(X)$ with $\|B\| < \delta$;
- (b) $\alpha(T - \lambda I)$ is a constant $\leq \alpha(T)$, $\beta(T - \lambda I)$ is a constant $\leq \beta(T)$ for $0 < |\lambda| < \delta$.

The above theorem shows that $\mathcal{A}(X)$ and $\mathcal{SF}(X)$ are open subsets of $\mathcal{L}(X)$. Furthermore, the continuity of the index shows that the *jump* of $T \in \mathcal{SF}(X)$

$$j(T) = \begin{cases} \alpha(T) - \alpha(T - \lambda I) & (0 < |\lambda| < \delta) \text{ if } \alpha(T) < \infty \\ \beta(T) - \beta(T - \lambda I) & (0 < |\lambda| < \delta) \text{ if } \beta(T) < \infty \end{cases}$$

is unambiguously defined.

PROPOSITION 1. *If $T \in \mathcal{SF}(X)$ then $j(T) = 0 \iff N(T) \subseteq \bigcap_{n \geq 1} T^n(X)$.*

PROOF. [14, Proposition 2.2]. □

We now list various classes of bounded linear operators which will be discussed:

$$\mathcal{SF}_0(X) = \{T \in \mathcal{SF}(X) : \alpha(T) = 0 \text{ or } \beta(T) = 0\};$$

$$\mathcal{B}(X) = \{T \in \mathcal{L}(X) : N(T) \subseteq T(X)\};$$

$$\mathcal{M}(X) = \{T \in \mathcal{L}(X) : T \text{ is left or right invertible in } \mathcal{L}(X)\};$$

$$\mathcal{D}(X) = \{T \in \mathcal{C}(X) : N(T) \subseteq \bigcap_{n \geq 1} T^n(X)\};$$

$$\mathcal{S}(X) = \{T \in \mathcal{R}(X) : N(T) \subseteq \bigcap_{n \geq 1} T^n(X)\}.$$

It is well known that $\mathcal{M}(X)$ is open. $\mathcal{SF}_0(X)$ is open by Theorem 1. An operator in $\mathcal{S}(X)$ is called an operator of Saphar type. Such operators have an important property:

$T \in \mathcal{S}(X)$ if and only if there is a neighbourhood $U \subset \mathbb{C}$ of 0 and a holomorphic function $F : U \rightarrow \mathcal{L}(X)$ such that

$$(T - \lambda I)F(\lambda)(T - \lambda I) = T - \lambda I \text{ for all } \lambda \in U.$$

For a proof see [7, Théorème 2.6] or [12, Theorem 1.4].

2 – Interior points of $\mathcal{D}(X)$ and $\mathcal{S}(X)$

If \mathcal{H} is a subset of $\mathcal{L}(X)$ we write $\text{int}(\mathcal{H})$ for the set of interior points of \mathcal{H} .

PROPOSITION 2. *If $T \in \text{int}(\mathcal{B}(X))$ then $N(T) = \{0\}$ or $T(X) = X$.*

PROOF. There exists $\delta > 0$ such that

$$S \in \mathcal{B}(X) \text{ whenever } \|T - S\| < \delta.$$

Suppose that $N(T) \neq \{0\}$ and $T(X) \neq X$. Then there are $x_0, y_0 \in X$ with $x_0 \neq 0$, $Tx_0 = 0$, $y_0 \notin T(X)$ and $\|Ty_0\| = \delta/2$. Since $y_0 \notin T(X)$ and $N(T) \subseteq T(X)$, we have $y_0 \notin N(T)$. An application of the Hahn-Banach extension theorem shows the existence of a continuous linear functional f such that

$$\alpha = f(x_0) \neq 0, \quad f(y_0) = 0 \quad \text{and} \quad \|f\| = 1$$

(see [6, Satz 36.3]). Define $S \in \mathcal{L}(X)$ by

$$Sx = Tx + f(x)Ty_0 \quad (x \in X).$$

It follows that $\|Tx - Sx\| = |f(x)|\|Ty_0\| \leq \|x\|\delta/2$, thus $\|T - S\| < \delta$, hence $S \in \mathcal{B}(X)$. Since $S(X) \subseteq T(X)$, we conclude that

$$N(S) \subseteq T(X).$$

Now put $z = y_0 - x_0/\alpha$. It results that

$$Sz = Ty_0 + f\left(y_0 - \frac{1}{\alpha}x_0\right)Ty_0 = Ty_0 - Ty_0 = 0.$$

This gives $z \in T(X)$, hence $y_0 = z + x_0/\alpha \in T(X) + N(T) = T(X)$ which contradicts $y_0 \notin T(X)$. \square

It is shown in [10] that neither $\mathcal{D}(X)$ nor $\mathcal{S}(X)$ are open subsets of $\mathcal{L}(X)$. But the following perturbation results are valid:

Suppose $T \in \mathcal{S}(X)$ (resp. $T \in \mathcal{D}(X)$), $B \in \mathcal{L}(X)$ and $B \left(\bigcap_{n=1}^{\infty} T^n(X) \right) \subseteq \bigcap_{n=1}^{\infty} T^n(X)$. If $\|B\|$ is sufficiently small then $T - B \in \mathcal{S}(X)$ (resp. $T - B \in \mathcal{D}(X)$).

(For proofs see [1, p. 150] (resp. [10, Corollaire 3.6]).)

Therefore a natural question arises: What are the interior points of $\mathcal{S}(X)$ and $\mathcal{D}(X)$? The following result gives an answer.

THEOREM 2.

- (a) $\text{int}(\mathcal{D}(X)) = \text{int}(\mathcal{B}(X) \cap \mathcal{C}(X)) = \mathcal{SF}_0(X)$.
 (b) $\text{int}(\mathcal{S}(X)) = \text{int}(\mathcal{B}(X) \cap \mathcal{R}(X)) = \mathcal{M}(X)$.

PROOF. (a) By Theorem 1 and Proposition 1, $\mathcal{SF}_0(X) \subseteq \mathcal{D}(X)$. Since $\mathcal{SF}_0(X)$ is open and $\mathcal{SF}_0(X) \subseteq \mathcal{D}(X) \subseteq \mathcal{B}(X) \cap \mathcal{C}(X)$, we have

$$\mathcal{SF}_0(X) \subseteq \text{int}(\mathcal{D}(X)) \subseteq \text{int}(\mathcal{B}(X) \cap \mathcal{C}(X)).$$

If $T \in \text{int}(\mathcal{B}(X) \cap \mathcal{C}(X))$ then $T \in \text{int}(\mathcal{B}(X))$, thus $\alpha(T) = 0$ or $\beta(T) = 0$, by Proposition 2. Since $T(X)$ is closed, we derive $T \in \mathcal{SF}_0(X)$.

(b) Since $\mathcal{M}(X)$ is open and $\mathcal{M}(X) \subseteq \mathcal{S}(X) \subseteq \mathcal{B}(X) \cap \mathcal{R}(X)$, we have

$$\mathcal{M}(X) \subseteq \text{int}(\mathcal{S}(X)) \subseteq \text{int}(\mathcal{B}(X) \cap \mathcal{R}(X)).$$

Let $T \in \text{int}(\mathcal{B}(X) \cap \mathcal{R}(X))$. There is $S \in \mathcal{L}(X)$ with $TST = T$. Proposition 2 shows that $(I - ST)(X) = N(T) = \{0\}$ or $TS(X) = T(X) = X$, thus $ST = I$ or $TS = I$, therefore $T \in \mathcal{M}(X)$. \square

REMARK. If X is a Hilbert space, then $\mathcal{C}(X) = \mathcal{R}(X)$ [1, p. 12], hence $\mathcal{D}(X) = \mathcal{S}(X)$. In this special case it was shown in [8, Théorème 6.5] that $\text{int}(\mathcal{D}(X)) = \mathcal{M}(X)$.

COROLLARY 1. *If X is a Hilbert space then $\text{int}(\mathcal{S}(X))$ is dense in $\mathcal{L}(X)$.*

PROOF. $\mathcal{M}(X)$ is dense in $\mathcal{L}(X)$ [4, Problem 140]. Now use Theorem 2. \square

COROLLARY 2.

- (a) $\text{int}(\{T \in \mathcal{SF}(X) : j(T) = 0\}) = \mathcal{SF}_0(X)$.
 (b) $\text{int}(\{T \in \mathcal{A}(X) : j(T) = 0\}) = \mathcal{M}(X)$.

PROOF. (a) follows from $\mathcal{SF}_0(X) \subseteq \{T \in \mathcal{SF}(X) : j(T) = 0\} \subseteq \mathcal{D}(X)$ (Proposition 1) and from Theorem 2.

(b) follows from $\mathcal{M}(X) \subseteq \{T \in \mathcal{A}(X) : j(T) = 0\} \subseteq \mathcal{S}(X)$ and from Theorem 2. \square

3 – The reduced minimum modulus of operators in $\mathcal{D}(X)$

By definition, the reduced minimum modulus $\gamma(T)$ of $T \in \mathcal{L}(X) \setminus \{0\}$ is given by

$$\gamma(T) = \inf \left\{ \frac{\|Tx\|}{d(x, N(T))} : x \in X, Tx \neq 0 \right\}.$$

($d(x, N(T))$ denotes the distance of x to $N(T)$.) Observe that $\gamma(T) > 0$ if and only if $T \in \mathcal{C}(X)$ [3, Theorem IV. 1.6].

PROPOSITION 3. *Let $T \in \mathcal{L}(X)$.*

- (a) *If $T \in \mathcal{D}(X)$ then $T^n \in \mathcal{D}(X)$ for all $n \in \mathbb{N}$.*
- (b) *If $T \in \mathcal{D}(X)$ then $\gamma(T^{n+m}) \geq \gamma(T^n)\gamma(T^m)$ for all $n, m \in \mathbb{N}$.*
- (c) *If $T \in \mathcal{R}(X)$ and $TST = T$ for some $S \in \mathcal{L}(X)$ then $\|S\|^{-1} \leq \gamma(T)$.*
- (d) *If $T \in \mathcal{S}(X)$ and $TST = T$ for some $S \in \mathcal{L}(X)$ then $T^n S^n T^n = T^n$ for each $n \in \mathbb{N}$.*

PROOF. (a) [11, Satz 6]. (b) [2, Lemma 1]. (c) [2, Lemma 4]. (d) [13, Proposition 2]. \square

We denote by $\sigma(T)$ the spectrum of $T \in \mathcal{L}(X)$ and by $r(T) = \max\{|\lambda| : \lambda \in \sigma(T)\}$ ($= \lim_{n \rightarrow \infty} \|T^n\|^{1/n}$) the spectral radius of T . $\partial\sigma(T)$ denotes the boundary of $\sigma(T)$.

PROPOSITION 4. *Let $T \in \mathcal{L}(X)$.*

- (a) *If $\mu \in \partial\sigma(T)$ then $T - \mu I \notin \mathcal{D}(X)$.*
- (b) *If $T \in \mathcal{D}(X)$ then*

$$\sup_{n \geq 1} \gamma(T^n)^{1/n} \leq \min \{|\mu| : \mu \in \partial\sigma(T)\},$$

the sequence $(\gamma(T^n)^{1/n})_{n \geq 1}$ converges and

$$\lim_{n \rightarrow \infty} \gamma(T^n)^{1/n} = \sup_{n \geq 1} \gamma(T^n)^{1/n}.$$

- (c) *If $T \in \mathcal{S}(X)$ and $TST = T$ for some $S \in \mathcal{L}(X)$ then*

$$r(S)^{-1} \leq \lim_{n \rightarrow \infty} \gamma(T^n)^{1/n}.$$

PROOF. (a) follows from [11, Satz 2].

(b) Fix $\mu \in \partial\sigma(T)$ such that $|\mu| = \min\{|\lambda| : \lambda \in \partial\sigma(T)\}$ and suppose that $|\mu| < \gamma(T^m)^{1/m}$ for some $m \in \mathbb{N}$. Thus $|\mu^m| < \gamma(T^m)$. Since $T^m \in \mathcal{D}(X)$ (Proposition 3(a)), Théorème 2.10 in [9] gives $T^m - \mu^m I \in \mathcal{D}(X)$. [11, Satz 6] implies now that $T - \mu I \in \mathcal{D}(X)$, but this contradicts (a). Hence $\gamma(T^m)^{1/m} \leq |\mu|$ for each $m \in \mathbb{N}$.

(b) follows from [2, remarks in connection with Lemma 1].

(c) By Proposition 3(d), $T^n S^n T^n = T^n$ for all $n \in \mathbb{N}$. Part (c) of Proposition 3 implies that $\|S^n\|^{-1} \leq \gamma(T^n)$ for each $n \in \mathbb{N}$, hence

$$r(S)^{-1} = \lim_{n \rightarrow \infty} \frac{1}{\|S^n\|^{1/n}} \leq \lim_{n \rightarrow \infty} \gamma(T^n)^{1/n}.$$

□

The following theorem is another perturbation result for operators in $\mathcal{D}(X)$ which generalizes Théorème 2.10 in [9].

THEOREM 3. *If $T \in \mathcal{D}(X)$, $B \in \mathcal{L}(X)$, $TB = BT$ and $r(B) < \lim_{n \rightarrow \infty} \gamma(T^n)^{1/n}$, then $T - B \in \mathcal{D}(X)$.*

PROOF. Since $r(B) = \inf_{k \geq 1} \|B^k\|^{1/k} < \sup_{n \geq 1} \gamma(T^n)^{1/n}$, there exists $k \in \mathbb{N}$ such that $\|B^{k+1}\| < \gamma(T^{k+1})$. By Proposition 3(a), $T^{k+1} \in \mathcal{D}(X)$, thus $T^{k+1} - B^{k+1} \in \mathcal{D}(X)$, by [10, Corollaire 3.6], since $B \left(\bigcap_{n=1}^{\infty} T^n(X) \right) \subseteq \bigcap_{n=1}^{\infty} T^n(X)$. $TB = BT$ implies

$$T^{k+1} - B^{k+1} = (T - B)(T^k + T^{k-1}B + \cdots + TB^{k-1} + B^k).$$

Therefore [11, Satz 5] shows that $T - B \in \mathcal{D}(X)$. □

The next result is proved in [10, Théorème 3.7]. It is now an immediate consequence of the last theorem.

THEOREM 4. *Let $T, Q \in \mathcal{L}(X)$. If Q is quasi-nilpotent and commutes with T , then*

$$T \in \mathcal{D}(X) \text{ if and only if } T - Q \in \mathcal{D}(X).$$

We close this paper with a perturbation result concerning operators in $\mathcal{S}(X)$. For the proof we need the following proposition.

PROPOSITION 5. *If $A, B \in \mathcal{L}(X)$ commute and $AB \in \mathcal{S}(X)$, then $A, B \in \mathcal{S}(X)$.*

PROOF. [5, Theorem 10]. □

THEOREM 5. *Let $T, Q \in \mathcal{L}(X)$. If Q is quasi-nilpotent and commutes with T , then*

$$T \in \mathcal{S}(X) \text{ if and only if } T - Q \in \mathcal{S}(X).$$

PROOF. It suffices to prove the implication $T \in \mathcal{S}(X) \implies T - Q \in \mathcal{S}(X)$. Put $S \in \mathcal{L}(X)$ such that $TST = T$. By Proposition 3(a),(d), $T^n \in \mathcal{S}(X)$ and $T^n S^n T^n = T^n$ for each $n \in \mathbb{N}$. Put $S_n := S^n T^n S^n$ ($n \in \mathbb{N}$). It follows that $T^n S_n T^n = T^n$, $S_n T^n S_n = S_n$ and $\|S_n\|^{1/n} \leq \|S\|^2 \|T\|$. There exists $k \in \mathbb{N}$ such that $\|Q^{k+1}\|^{1/(k+1)} < (\|S\|^2 \|T\|)^{-1}$, thus $\|Q^{k+1}\| < \|S_{k+1}\|^{-1}$. By [1, Theorem 9 in Section 5.2], $T^{k+1} - Q^{k+1} \in \mathcal{S}(X)$. $TQ = QT$ implies

$$T^{k+1} - Q^{k+1} = (T - Q)(T^k + T^{k-1}Q + \dots + TQ^{k-1} + Q^k),$$

hence $T - Q \in \mathcal{S}(X)$, by Proposition 5. □

REFERENCES

- [1] S.R. CARADUS: *Generalized Inverses and Operator Theory*, Queen's Papers in Pure and Applied Math. No 50 (1978).
- [2] K.H. FÖRSTER – M.A. KAASHOEK: *The asymptotic behaviour of the reduced minimum modulus of a Fredholm operator*, Proc. Amer. Math. Soc. **49**, 123-131, (1975).
- [3] S. GOLDBERG: *Unbounded linear operators*, New York (1966).
- [4] P.R. HALMOS: *A Hilbert space problem book*, 2nd ed. Princeton (1980).

-
- [5] R. HARTE: *Taylor exactness and Kaplansky's lemma*, J. Operator Theory **25**, 399-416, (1991).
- [6] H. HEUSER: *Funktionalanalysis*, 2nd ed. Stuttgart (1986).
- [7] M. MBEKTHA: *Généralisation de la décomposition de Kato aux opérateurs paranormaux et spectraux*, Glasgow Math. J. **29**, 159-175, (1987).
- [8] M. MBEKTHA: *Résolvant généralisé et théorie spectrale*, J. Operator Theory **21**, 69-105, (1989).
- [9] M. MBEKTHA – A. OUAHAB: *Opérateurs s-régulier dans une espace de Banach et théorie spectrale*, Pub. IRMA, Lille Vol. 22, N° XII (1990).
- [10] M. MBEKTHA – A. OUAHAB: *Perturbations des opérateurs s-réguliers et continuité de certain sous-espaces dans le domaine quasi-Fredholm*, Pub. IRMA, Lille Vol. 24, N° X (1991).
- [11] CH. SCHMOEGER: *Ein Spektralabbildungssatz*, Arch. Math. **55**, 484-489, (1990).
- [12] CH. SCHMOEGER: *The punctured neighbourhood theorem in Banach algebras*, Proc. R. Ir. Acad. **91A**, No. 2, 205-218, (1991).
- [13] CH. SCHMOEGER: *Relatively regular operators and a spectral mapping theorem*, J. Math. Anal. Appl. **175**, 315-320, (1993).
- [14] T.T. WEST: *A Riesz-Schauder theorem for semi-Fredholm operators*, Proc. R. Ir. Acad. **87A**, No. 2, 137-146, (1987).

*Lavoro pervenuto alla redazione il 7 gennaio 1994
ed accettato per la pubblicazione il 13 aprile 1994*

INDIRIZZO DELL'AUTORE:

Christoph Schmoeger – Mathematisches Institut I – Universität Karlsruhe – Postfach 6980 –
76128 Karlsruhe 1 – Germany