

Extremal numbers for cycles in a hypercube

Maria Axenovich *

Abstract

Let $\text{ex}(Q_n, H)$ be the largest number of edges in a subgraph G of a hypercube Q_n such that there is no subgraph of G isomorphic to H . We show that for any integer $k \geq 3$

$$\text{ex}(Q_n, C_{4k+2}) = O(n^{\frac{5}{6} + \frac{1}{3(2k-2)}} 2^n).$$

A *hypercube* Q_n , where n is a natural number, is a graph on a vertex set $\{A : A \subseteq [n]\}$ and an edge set consisting of all pairs $\{A, B\}$, where $A \subseteq B$ and $|A| = |B| - 1$. Here, $[n] = \{1, \dots, n\}$. For a graph H , let $\text{ex}(Q_n, H)$ be the largest number of edges in a subgraph G of a hypercube Q_n such that there is no subgraph of G isomorphic to H . The behaviour of the function $\text{ex}(Q_n, H)$ is not well understood in general and it is not even known for what graphs H , $\text{ex}(Q_n, H) = \Omega(\|Q_n\|)$, where $\|Q_n\| = n2^{n-1}$ denotes the number of edges of Q_n . See the following papers on the topic: [1–5, 16, 17]. Conlon [10] observed a connection between extremal numbers in the hypercube and classical extremal numbers for uniform hypergraphs. That allowed to determine a large class of graphs H such that $\text{ex}(Q_n, H) = o(\|Q_n\|)$, i.e., the class of graphs with zero Turán density.

In this note we focus on the case when H is an even cycle. Here, a 2ℓ -cycle, denoted $C_{2\ell}$, is a graph on 2ℓ vertices and 2ℓ edges such that the edges are pairs of consecutive vertices with respect to some circular ordering of the vertices. It is known that $\text{ex}(Q_n, C_4) = \Omega(\|Q_n\|)$ and $\text{ex}(Q_n, C_6) = \Omega(\|Q_n\|)$, see Chung [8], Conder [9], and Brass et al. [6]. Chung [8] showed that $\text{ex}(Q_n, C_{4k}) = o(\|Q_n\|)$, for any integer $k \geq 2$. Füredi and Özkahya [14, 15] extended Chung's results by showing that $\text{ex}(Q_n, C_{4k+2}) = o(\|Q_n\|)$, for any integer $k \geq 3$. Thus if $\ell \notin \{2, 3, 5\}$, $\text{ex}(Q_n, C_{2\ell}) = o(\|Q_n\|)$. While $\text{ex}(Q_n, C_{2\ell}) = \Omega(\|Q_n\|)$ for $\ell \in \{2, 3\}$, it remains unclear whether $\text{ex}(Q_n, C_{10}) = \Omega(\|Q_n\|)$.

Conlon [10] noted that one can obtain a non-trivial lower bound on $\text{ex}(Q_n, C_{2\ell})$ using Lovász Local Lemma. For $\ell \geq 2$, $\text{ex}(Q_n, C_{2\ell}) = \Omega(n^{\frac{1}{2} + \frac{1}{4\ell-2}} 2^n)$, as proven using deletion method, see Conlon's homepage [11]. In Appendix, we include the proof of this lower bound using Lovász Local Lemma.

Considering more specific upper bounds for cycles with zero Turán density, Conlon [10] proved for $k \geq 2$ that $\text{ex}(Q_n, C_{4k}) \leq c_k n^{-\frac{1}{2} + \frac{1}{2k}} \|Q_n\|$. Füredi and Özkahya [15], [14] showed that $\text{ex}(Q_n, C_{4k+2}) = O(n^{-q_k} \|Q_n\|)$, where $q_k = \frac{1}{2k+1}$ for $k \in \{3, 5, 7\}$, and $q_k = \frac{1}{16} - \frac{1}{16(k-1)}$ for any other integer $k \geq 3$. Note that for large k , this upper bound on $\text{ex}(Q_n, C_{4k+2})$ behaves as $n^{\frac{15}{16} + \epsilon} 2^n$, for small positive ϵ . Here, we improve the leading term in the exponent from $\frac{15}{16}$ to $\frac{5}{6}$:

Theorem 1. *Let ℓ be an odd integer, $\ell \geq 7$. Then $\text{ex}(Q_n, C_{2\ell}) = O\left(n^{\frac{5}{6} + \frac{1}{3(\ell-3)}} 2^n\right)$.*

*Karlsruhe Institute of Technology, Karlsruhe, Germany; email: maria.aksenovich@kit.edu.

The main ingredient of the proof is also due to Conlon [10] using partite representation. Our contribution here is finding a suitable representation and bounding the respective extremal hypergraph function.

We need some definitions and notations. For a set V and a non-negative integer r , we denote by $\binom{V}{r}$ the set of all r -element subsets of V . An r -graph on a vertex set V is a pair (V, E) , where $E \subseteq \binom{V}{r}$ is the set of edges. Note that a graph is a 2-graph. We denote the number of edges in an r -graph H by $\|H\|$. If H is an r -graph, then $\text{ex}(n, H)$ denotes the largest number of edges in an n -vertex r -graph not containing H as a subgraph. An r -graph is k -partite if its vertex set could be partitioned into k parts such that any edge contains at most one vertex from each part. Edges $\{x, y\}$ and $\{x, y, z\}$ are written simply as xy and xyz , respectively. For a 3-graph \mathcal{H} and its vertex x , the *link graph* $L(x)$ is the graph with no isolated vertices whose edge set consists of all pairs yz such that xyz is an edge of \mathcal{H} . Let H be a bipartite graph. Let the *two-lift* of H be the 3-graph \mathcal{H} with vertex set $\{a, b\} \cup V(H)$, where $a \notin V(H)$ and $b \notin V(H)$, $a \neq b$, each edge of \mathcal{H} contains exactly one of a or b , and $L(a) = L(b) = H$.

For a non-negative k , the k th layer of Q_n is the set $\binom{[n]}{k}$ of vertices. We say that a subgraph H of Q_n has a k -partite representation \mathcal{H} if H is isomorphic to a graph H' with a vertex set contained in $\binom{[n]}{k-1} \cup \binom{[n]}{k}$ such that $V(H') \cap \binom{[n]}{k}$ is an edge-set of a k -partite k -graph. In this case, we denote such a hypergraph with no isolated vertices by $\mathcal{H}(H)$ and call it a k -partite representation of H . For example, if H is an 8-cycle, it has a 2-partite representation with edges 12, 23, 34, 14 corresponding to an 8-cycle with vertices 1, 12, 2, 23, 3, 34, 4, 14, 1, in order. A more general result by Conlon implies the following in the case when $k = 3$.

Theorem 2 (Conlon [10]). *If H is a subgraph of Q_n that has a 3-partite representation \mathcal{H} such that $\text{ex}(n, \mathcal{H}) \leq \alpha n^3$, then $\text{ex}(Q_n, H) \leq \alpha^{1/3} 2^n n$.*

Lemma 1. *A cycle $C_{2\ell}$, where $\ell \geq 7$ is an odd integer, has a 3-partite representation \mathcal{H}_ℓ such that \mathcal{H}_ℓ is a subgraph of a two-lift of a cycle of length $\ell - 3$ with a pendant edge.*

Proof. We shall construct a cycle $C = C_{2\ell}$ for odd $\ell = 2k + 1$ with vertices in the third and second layer of Q_n . We shall define A_ℓ to be the ordered set of vertices of C in the third layer, B_C to be a respective set of vertices in the second layer, and let C be built by taking vertices from A_ℓ and B_ℓ alternately in order. Let

$$\begin{aligned} A_{2k+1} &= (ax_1y_1, ax_2y_1, ax_2y_2, \dots, ax_{k-1}y_{k-2}, ax_{k-1}y_{k-1}, bx_{k-1}y_{k-1}, bx_{k-1}y_0, bx_1y_0, bx_1y_1), \\ B_{2k+1} &= (ay_1, ax_2, ay_2, \dots, ax_{k-1}, x_{k-1}y_{k-1}, bx_{k-1}, by_0, bx_1, x_1y_1). \end{aligned}$$

Then C has a 3-partite representation \mathcal{H}_ℓ with the edge set equal to the underlying unordered set of A_ℓ . Indeed, we see that H_ℓ is a three-partite 3-graph with parts $\{a, b\}$, $\{x_1, x_2, \dots, x_{k-1}\}$, and $\{y_0, y_1, y_2, \dots, y_{k-1}\}$. One can visualise H_ℓ by considering link graphs $L(a)$ and $L(b)$ - these graphs are paths of lengths $2k - 3$ and 4 respectively, sharing only vertices x_1, y_1, x_{k-1} , and y_{k-1} and edges x_1y_1 and $x_{k-1}y_{k-1}$. The union of these two paths is a cycle of length $2(k - 1) = \ell - 3$ and a pendant edge. \square

Lemma 2. *Let H be a bipartite graph and \mathcal{H} be a two-lift of H . If $\text{ex}(n, H) = O(n^\gamma)$, then $\text{ex}(n, \mathcal{H}) = O(n^{\frac{\gamma+4}{2}})$.*

Proof. Let $\gamma, c' > 0$, n sufficiently large, and $\text{ex}(n, H) < c'n^\gamma$. Let \mathcal{G} be a 3-graph with vertex set V , $|V| = n$ and cn^σ edges, where $\sigma = \frac{\gamma+4}{2}$ and c is a constant, $c' = c^2$. We shall show that \mathcal{G} contains a two-lift \mathcal{H} of H .

Let $U = \binom{V}{2}$ and G be a bipartite graph with partite sets V and U and edge set $\{\{x, zz'\} : x \in V, zz' \in U, xzz' \in E(\mathcal{G})\}$. Then $\|G\| = 3\|\mathcal{G}\| = 3cn^\sigma$. Let q be the largest integer such that G contains a complete bipartite subgraph with two vertices in V and q vertices in U . We shall show that $q \geq n^{2\sigma-4}$.

To do this, we shall apply a standard double counting and reduction to smaller uniformity, already used by Erdős [13] in bounding the extremal function for complete k -partite k -graphs. Let X be the number of stars in G with 2 edges and center in U . Then $X \geq \sum_{u \in U} \binom{\deg_G(u)}{2} \geq |U| \binom{\|G\|/|U|}{2} \geq c^2 n^2 n^{2(\sigma-2)}$. On the other hand, since every pair of vertices from V belongs to at most q such stars, $X \leq q \binom{n}{2} \leq qn^2$. Thus $q \geq c^2 n^{2\sigma-4} = c'n^\gamma$. Let $Q \subseteq U$, $|Q| = q$, $a, b \in V$, such that a, b , and Q induce $K_{2,q}$ in G . Then a, b and Q correspond to a two-lift \mathcal{H}' of a graph H' with the edge-set Q . Since $|Q| = q = c'n^\gamma > \text{ex}(n, H)$, we see that H is a subgraph of H' . Thus \mathcal{G} contains a two-lift of H . \square

Proof of Theorem 1. It was shown by Erdős [12] as well as by Bondy and Simonovits [7], that $\text{ex}(n, C_{2m}) = O(n^{1+\frac{1}{m}})$. This implies that $\text{ex}(n, H) \leq O(n^{1+\frac{2}{\ell-3}})$, where H is a cycle of length $\ell - 3$ with a pendant edge, for odd $\ell \geq 7$. Now, using Theorem 2 and Lemmas 1 and 2, with $\gamma = 1 + \frac{2}{\ell-3}$, $\frac{\gamma+4}{2} = \frac{5+\frac{2}{\ell-3}}{2}$, and $\alpha = -\frac{1}{2} + \frac{1}{\ell-3}$, we have $\text{ex}(Q_n, C_{2\ell}) \leq O(\alpha^{1/3} n^{2^n}) = O(n^{1-\frac{1}{6}+\frac{1}{3(\ell-3)} 2^n})$. \square

Remark After uploading this note, the author learned about a better bound obtained by Istvan Tomon [18], page 28., as a corollary of a much more general result on topologically defined hypergraphs: $\text{ex}(Q_n, C_{2\ell}) = O(n^{\frac{2}{3}+\delta} 2^n)$, for some $\delta = O(\frac{\log \ell}{\ell})$.

Acknowledgements The author thanks David Conlon for bringing her attention to [11], Frithjof Marquardt for discussions, and Zixiang Xu for pointing out [18]. The research was supported in part by the DFG grant FKZ AX 93/2-1.

References

- [1] N. Alon, A. Krech and T. Szabó, Turán's theorem in the hypercube. *SIAM Journal on Discrete Mathematics*, 21(1) (2007), 66–72.
- [2] N. Alon, R. Radoičić, B. Sudakov and J. Vondrák, A Ramsey-type result for the hypercube. *J. Graph Theory* 53 (2006), no. 3, 196–208.
- [3] M. Axenovich and R. Martin, A note on short cycles in the hypercube. *Discrete Math.* 306 (2006), 2212–2218.
- [4] R. Baber, Turán densities of hypercubes. *ArXiv preprint* (2012) arXiv:1201.3587.
- [5] J. Balogh, P. Hu, B. Lidický and H. Liu, Upper bounds on the size of 4- and 6-cycle-free subgraphs of the hypercube. *European Journal of Combinatorics* 35 (2014), 75–85.
- [6] P. Brass, H. Harborth and H. Nienborg, On the maximum number of edges in a C_4 -free subgraph of Q_n . *Journal of Graph Theory* 19, 17–23.
- [7] J. A. Bondy and M. Simonovits, Cycles or even length in graphs, *J. Combin. Theory Ser. B* 16 (1974), 97–105.
- [8] F. Chung, Subgraphs of a hypercube containing no small even cycles. *Journal Graph Theory* 16 (1992), 273–286.
- [9] M. Conder, Hexagon-free subgraphs of hypercubes. *Journal of Graph Theory* 17 (4) (1993), 477–479.
- [10] D. Conlon, An extremal theorem in the hypercube. *Electronic Journal of Combinatorics* 17 (2010).
- [11] D. Conlon, <http://www.its.caltech.edu/~dconlon/HyperCycleErratum.pdf>
- [12] P. Erdős, unpublished
- [13] P. Erdős, On extremal problems of graphs and generalized graphs. *Israel J. Math.* 2 (1964) 183–190.

- [14] Z. Füredi and L. Özkahya, On even-cycle-free subgraphs of the hypercube. *Electronic Notes in Discrete Mathematics* 34 (2009), 515–517.
- [15] Z. Füredi and L. Özkahya, On even-cycle-free subgraphs of the hypercube. *Journal of Combinatorial Theory, Series A*, 118 (2011), 1816–1819.
- [16] D. Offner, Some Turán type results on the hypercube. *Discrete Mathematics*, 309(9) (2009), 2905–2912.
- [17] A. Thomason and P. Wagner, Bounding the size of square-free subgraphs of the hypercube. *Discrete Math.* 309 (2009), 1730–1735.
- [18] I. Tomon, Robust (rainbow) subdivisions and simplicial cycles. ArXiv preprint 2022: arXiv:2201.12309.

1 Appendix

Theorem 3. For any integer ℓ , $\ell \geq 2$, $ex(Q_n, C_{2\ell}) = \Omega(n^{\frac{1}{2} + \frac{1}{4\ell-2}} 2^n)$.

Proof. Consider a random edge-coloring of Q_n , where the colors for the edges are chosen i.i.d. with probability $p = cn^{-a}$ for each of the $1/p$ colors, where the constants c and a will be determined later. For a copy C or $C_{2\ell}$, let A_C be the probability of a bad event that all edges of C have the same color. Denoting $P = \text{Prob}(A_C)$, we have $P \leq p^{2\ell-1}$. Note that A_C is mutually independent of all events $A_{C'}$ except for those C' 's that share an edge with C .

To determine the maximum degree in the dependency graph, consider first $N(Q_n, C_{2\ell})$, the number of 2ℓ -cycles in Q_n . Represent the vertices of Q_n as binary vectors, consider a 2ℓ -cycle C and the set S_C of all positions at which some two adjacent in C vertices differ. Then $|S_C| \leq \ell$. Since there are at most n^ℓ ways to choose the set S_C and at most 2^n ways to fix the remaining positions, in which all vertices of C coincide, we have that $N(Q_n, C_{2\ell}) \leq c_\ell n^\ell 2^n$.

Let x be the number of 2ℓ -cycles containing a given edge. By symmetry of Q_n we see that x is independent of the choice of the edge. Let X be the set of pairs (C, e) , where C is a cycle of length 2ℓ in Q_n and e is an edge of C . Then $|X| = N(Q_n, C_{2\ell}) \cdot 2\ell$ and $|X| = n2^{n-1} \cdot x$, implying that $x = N(Q_n, C_{2\ell}) \cdot 2\ell / (n2^{n-1}) \leq c_\ell n^\ell 2^n 2\ell / (n2^{n-1}) = c'_\ell n^{\ell-1}$. Thus the maximum degree D of the dependency graph satisfies $D \leq 2\ell x \leq 2\ell \cdot c'_\ell n^{\ell-1}$.

Observe that $P(D+1)e = (cn^{-a})^{2\ell-1}(c'_\ell n^{\ell-1} + 1)e$, it is less than 1 if the constant c in the definition of p is sufficiently small and the exponent $-a(2\ell-1) + \ell - 1 = 0$, i.e., if $a = (\ell-1)/(2\ell-1)$. Using Lovász Local Lemma, we have that with a positive probability there are no bad events A_C . A largest color class gives a $C_{2\ell}$ -free subgraph of Q_n with at least $p||Q_n|| = cn^{1-a}2^n = cn^{\frac{\ell}{2\ell-1}}2^n = cn^{\frac{1}{2} + \frac{1}{4\ell-2}}2^n$ edges. \square