

A class of graphs of zero Turán density in a hypercube

Maria Axenovich *

Abstract

A graph is cubical if it is a subgraph of a hypercube. For a cubical graph H and a hypercube Q_n , $\text{ex}(Q_n, H)$ is the largest number of edges in an H -free subgraph of Q_n . If $\text{ex}(Q_n, H)$ is at least a positive proportion of the number of edges in Q_n , H is said to have a positive Turán density in a hypercube or simply a positive Turán density; otherwise it has a zero Turán density. Determining $\text{ex}(Q_n, H)$ and even identifying whether H has a positive or a zero Turán density remains a widely open question for general H . By relating extremal numbers in a hypercube and certain corresponding hypergraphs, Conlon found a large class of cubical graphs, ones having so-called partite representation, that have a zero Turán density. He raised a question whether this gives a characterisation, i.e., whether a cubical graph has zero Turán density if and only if it has partite representation. Here, we show that, as suspected by Conlon, this is not the case. We give an example of a class of cubical graphs which have no partite representation, but on the other hand, have a zero Turán density. In addition, we show that any graph whose every block has partite representation has a zero Turán density in a hypercube.

1 Introduction

A *hypercube* Q_n with a ground set X of size n , is a graph on a vertex set $\{A : A \subseteq X\}$ and an edge set consisting of all pairs $\{A, B\}$, where $A \subseteq B$ and $|A| = |B| - 1$. Unless specified, $X = [n]$, where $[n] = \{1, \dots, n\}$. We often identify vertices of Q_n with binary vectors that are indicator vectors of respective sets. If a graph is a subgraph of Q_n , for some n , it is called *cubical*. We denote the number of vertices and the number of edges in a graph H by $|H|$ and $\|H\|$, respectively. The degree of a vertex y in a graph H is denoted $d(y)$ or $d_H(y)$. A *block* in a graph is a maximal two-connected subgraph or a bridge. We shall need the notion of layers. The i th *vertex layer* of Q_n , denoted V_i , is the set of vertices $\binom{[n]}{i}$, $i = 0, \dots, n$. The i th *edge-layer* L_i of Q_n is a graph induced by $V_i \cup V_{i-1}$, $i = 1, \dots, n$.

For a graph H , let the *extremal number* of H in Q_n , denoted $\text{ex}(Q_n, H)$, be the largest number of edges in a subgraph G of Q_n such that there is no subgraph of G isomorphic to H . A graph H is said to have *zero Turán density in a hypercube* if $\text{ex}(Q_n, H) = o(\|Q_n\|)$. Otherwise, we say that H has a *positive Turán density in a hypercube*. Note that by using a standard double counting argument, the sequence $\text{ex}(Q_n, H)/\|Q_n\|$ is non-increasing, thus the above density notions are well-defined. When clear from context, we simply say Turán density instead of Turán density in a hypercube. The behaviour of the function $\text{ex}(Q_n, H)$ is not well understood in general and it is not even known what graphs have positive or zero Turán density. Currently, the only known cubical graphs of positive Turán density are those containing a 4- or a 6-cycle as a subgraph, [8, 9], and one special graph of girth 8 [2]. Conlon [10] observed a connection between extremal numbers in a hypercube and classical extremal numbers for uniform hypergraphs. That permitted to determine large specific classes of graphs, such as for example subdivisions, with zero

*Karlsruhe Institute of Technology, Karlsruhe, Germany; email: maria.aksenovich@kit.edu

Turán density, see [2]. For more results on extremal numbers in a hypercube, see [1, 5, 6, 16, 17].

A graph H has a k -partite representation \mathcal{H} if for some n , H is isomorphic to a subgraph H' of the k th layer L_k of Q_n such that $V(H') \cap V_k$ is an edge-set of a k -partite k -uniform hypergraph \mathcal{H} . The graph H' is called *representing*. If H has a k -partite representation for some k , we say that H has a *partite representation*. Here, a k -uniform hypergraph is k -partite if the vertex set can be partitioned into k parts such that each hyperedge has exactly one vertex in each part. Here, we omit the brackets and commas denoting sets, when clear from context, i.e., for a set $\{1, 2\}$ we simply write 12. For example, if H is an 8-cycle, it has a 2-partite representation \mathcal{H} with hyperedges 12, 23, 34, 14 corresponding to an 8-cycle with vertices 1, 12, 2, 23, 3, 34, 4, 14, 1, in order. For a k -uniform hypergraph \mathcal{H} , $\text{ex}_k(n, \mathcal{H})$ denotes the largest number of edges in a k -uniform n -vertex hypergraph with no subgraph isomorphic to \mathcal{H} .

Using a theorem by Erdős [11], that states that $\text{ex}_k(n, \mathcal{H}) = o_n(n^k)$ for any k -partite hypergraph \mathcal{H} , Conlon [10] proved that if H is a graph that has partite representation, then $\text{ex}(Q_n, H) = o(\|Q_n\|)$. In the same paper Conlon [10] raised a question whether cubical graphs that have no partite representation have positive Turán density. Here, we show that it is not the case, i.e. having a partite representation is not a characterisation for zero Turán density in a hypercube. For that, we construct a family of cubical graphs that have no partite representation, but have zero Turán density. These graphs are formed by two copies of so-called *theta*-graphs that we define below.

For a graph G , let $G(1)$ be a 1-subdivision of G , i.e., a graph G' with a vertex set $V(G) \cup E(G)$ and an edge set $\{ue, ev : e = uv \in E(G)\}$. We call the vertices from $V(G)$ in G' , the *poles* of G' and other vertices, the *subdivision vertices*. Let $K_{s,t}$ denote a complete bipartite graph with parts of sizes s and t , respectively. We shall be considering the 1-subdivision of $K_{q,2}$, $q \geq 3$. We shall call the two vertices of degree q in $K_{q,2}(1)$, the *main poles*. Note that $K_{q,2}(1)$ is also referred to as a *theta graph* with q legs of length 4. Here the legs are paths with endpoints being main poles. We shall use a shorter notation $\Theta(q)$ for $K_{q,2}(1)$, when appropriate. If a graph G contains a subgraph isomorphic to H , we call such a subgraph a *copy of H* .

Let $H(q)$ be a union of two copies of $\Theta(q)$ sharing exactly one vertex that is a main pole of one copy and a subdivision vertex in another copy, see Figure 1.




Figure 1: A graph $H(3)$ with larger vertices being poles in the respective copies of theta graphs as well as a nice coloring of one theta graph

Marquardt [15] showed that $H(q)$ has no partite representation for $q = 3$. Here, we give a

slightly different proof of this fact for any $q \geq 3$ and show that $H(q)$ has a zero Turán density. In doing so, we prove a result of an independent interest that gives a new class of cubical graphs of zero Turán density.

Theorem 1. *Let H be a graph whose every block has a partite representation. Then H has a zero Turán density in a hypercube.*

Theorem 2. *For any $q \geq 3$ the graph $H(q)$ is cubical, has no partite representation, but has a zero Turán density in a hypercube.*

2 Proofs of main results

We shall need the following lemmas. The first one gives a property of graphs with partite representations, formulated by Marquardt [15], we include it here for completeness. The second one is basically a result of Conlon [10], but with a slightly different averaging argument that also gives a more precise statement about an embedding of a graph in a layer in two different ways. Finally, the last one is the main lemma needed for Theorem 1. If a graph H has a k -partite representation with representing graph H' , then the vertices of H corresponding to $V(H') \cap V_k$ are called *top vertices* with respect to this representation, all other vertices are called *bottom vertices*. Note that if a graph has a k -partite representation, it has a $(k + 1)$ -partite representation that could be seen by simply adding a new element to every vertex of a representing graph.

Lemma 3. *Let H and G be connected graphs, each having a k -partite representation. Assume that $V(H) \cap V(G) = \{v\}$, where v that is a top vertex for both H and G or a bottom vertex for both H and G . Then $H \cup G$ has a partite representation.*

Proof. Let H' and G' be copies of H and G that are subgraphs of the k th layer in hypercubes with a ground set X and Y , respectively, such that $V(H') \cap \binom{X}{k}$ and $V(H') \cap \binom{Y}{k}$ are the edge sets of k -partite hypergraphs with parts U_1, \dots, U_k and parts W_1, \dots, W_k , respectively. Let u and w be respective copies of v in H' and G' .

Assume first that v is a top vertex of H and of G . Assume that $X \cap Y = \emptyset$. Let F'' be an induced subgraph of a hypercube Q with ground set $X \cup Y$ and vertex set $\{x \cup w : x \in V(H')\} \cup \{u \cup y : y \in V(G')\}$. We see that F'' contains a copy F' of F and is contained in the layer $2k$ of Q with the vertex $u \cup w$ playing a role of v . Moreover $V(F') \cap \binom{X \cup Y}{2k}$ is an edge set of a $2k$ -partite hypergraph with parts $U_1, \dots, U_k, W_1, \dots, W_k$.

Assume now that v is a bottom vertex of H and of G . Assume further that $X \cap Y = [k - 1]$ and $u = w = [k - 1]$. Let F'' be an induced subgraph of a hypercube Q with ground set $X \cup Y$ and vertex set $V(H') \cup V(G')$. We see that F'' contains a copy F' of F and is contained in layer k of Q with vertex $v' = [k - 1]$ playing a role of v . Moreover $V(F') \cap \binom{X \cup Y}{k}$ is an edge set of a k -partite hypergraph with parts $U_1 \cup W_1, \dots, U_k \cup W_k$. \square

Lemma 4. *Let Z be a connected bipartite graph with a partite representation. Then for any $\gamma > 0$ there is $n_1 = n_1(\gamma, Z)$ such that for any $n > n_1$ the following holds. Let L_j be the j th layer of Q_n , where $n/2 - n^{2/3} \leq j \leq n/2 + n^{2/3}$. Let $G \subseteq L_j$ be a graph such that $\|G\| \geq \gamma \|L_j\|$. Fix one of the two partite set of Z arbitrarily and call its vertices odd. Then there is a copy of Z in G with odd vertices in V_j and there is a copy of Z in G with odd vertices in V_{j-1} .*

Proof. Let Z have a k -partite representation in a hypercube Q_n with ground set $[n]$, with odd vertices corresponding to k -element sets. We need the following notation. For any $x \in V(Q_n)$, let $\text{Up}(x)$ be the *up set* of x , i.e., $\text{Up}(x) = \{y \subseteq [n] : x \subseteq y\}$. Note that $\text{Up}(x)$ induces a graph isomorphic to Q_m , for $m = n - |x|$.

To prove the first part of the lemma, for each $x \in V_{j-k}$, let $U_x = U_{x,k}$ be the intersection of G and the k th edge-layer of the hypercube Q induced by $\text{Up}(x)$. Note that U_x is a subgraph of L_j , the j th layer of Q_n . We call a vertex y of U_x a *full vertex* if it is in the k th vertex layer of Q and has degree k in U_x , the largest possible degree. Let u_x be the number of full vertices in U_x . We shall argue that there is a vertex $x \in V_{j-k}$ such that u_x is large.

Let t be the number of k -edge stars in G with the center in V_j . Then

$$t = \sum_{y \in V_j} \binom{d_G(y)}{k} \geq |V_j| \binom{\gamma \|L_j\| / |V_j|}{k}.$$

Each such a star corresponds to a full vertex in U_x , for some x . Thus $\sum_{x \in V_{j-k}} u_x \geq t$ and there is a vertex x such that

$$u_x \geq \frac{|V_j|}{|V_{j-k}|} \binom{\gamma \|L_j\| / |V_j|}{k}.$$

Since $n/2 - n^{2/3} \leq j \leq n/2 + n^{2/3}$ and k is a fixed constant, we have that $\|L_j\| / |V_j| = \frac{n}{2}(1 + o(1))$ and $|V_j| / |V_{j-k}| \geq c'(k)$, so

$$u_x \geq c(k)n^k,$$

for positive constant $c(k)$ and $c'(k)$. Consider the full vertices in U_x . They correspond to u_x hyperedges in a k -uniform hypergraph with the vertex set $[n] - x$ of size $n - j + k = \frac{n}{2}(1 + o(1))$. By a theorem of Erdős [11], such a hypergraph contains any fixed k -partite k -uniform hypergraph, and thus in particular the one representing Z . Therefore G contains a copy of Z with odd vertices in V_j .

To prove the second part of the Lemma, we can repeat the above argument by considering the vertices x in V_{j+k-1} and their downsets. Alternatively, we see that L_j corresponds to $L_{j'}$, a ‘‘symmetric’’ layer, where $j' = n + 1 - j$, $V_{j'}$ corresponds to V_{j-1} , and $V_{j'-1}$ corresponds to V_j . Thus, finding a copy of Z in $L_{j'}$ with odd vertices in $V_{j'}$ corresponds to finding a copy of Z in L_j with odd vertices in V_{j-1} . Since j' satisfies the same conditions as j , i.e., $n/2 - n^{2/3} \leq j' \leq n/2 + n^{2/3}$, we thus could use the first part of the Claim to obtain the second one. \square

Lemma 5. *Let H be a connected bipartite graph with ℓ blocks, where every block has a partite representation. Then for any $\gamma > 0$ there is $n_0 = n_0(\gamma, H)$ such that for any $n > n_0$ the following holds. Let L_j be the j th layer of Q_n , where $n/2 - n^{2/3} \leq j \leq n/2 + n^{2/3}$. Let $G \subseteq L_j$ be a graph with $\|G\| = \gamma \|L_j\|$. Fix one of two partite sets of H arbitrarily and call its vertices odd. Then there is a copy of H in G with odd vertices in V_j and there is a copy of H in G with odd vertices in V_{j-1} .*

Proof. We shall prove the statement by induction on ℓ with the base case $\ell = 1$ directly following from Lemma 4.

If H has ℓ blocks, $\ell \geq 2$, such that each has a k -partite representation, let $H = H' \cup H''$, where H' and H'' share a single vertex v' , and H'' is a block of H , i.e., a leaf block. Let F be a

graph that is a union of $q > |V(H')|$ copies F_1, \dots, F_q of H'' that pairwise share only a vertex corresponding to v' . Let $\gamma > 0$ and n_0 be sufficiently large (we shall specify how large later). Let $G \subseteq L_j$, $\|G\| = \gamma\|L_j\|$.

Assume first that v' is an odd vertex.

The idea of the proof is as follows. We shall first find many copies of H' in G such that a vertex corresponding to v' is in V_j . Then we shall find a copy of F^* of F and a copy H^* of H' such that $V(F^*) \cap V(H^*) \cap V_j = \{v\}$, where v plays a role of v' . Finally, since q is large enough, we shall claim that there is a copy F_i^* of F_i for some i such that $V(F_i^*) \cap V(H^*) \cap V_{j-1} = \emptyset$. This will imply that $F_i^* \cup H^*$ is a copy of H . Next we shall give the details of this argument:

First, we shall construct sets \tilde{V}, V , and V' , such that $V' \subseteq V \subseteq \tilde{V} \subseteq V_j$ as follows:

Consider all copies H'_v of H' in G with a vertex playing a role of v' being in V_j . Let $\tilde{V} \subseteq V_j$ be the set of such vertices v . Note that $\|G[V_j - \tilde{V}, V_{j-1}]\| < \frac{1}{4}\gamma\|L_j\|$, otherwise by induction we can find another copy H'_w of H' with a vertex $w \in V_j - \tilde{V}$ playing a role of v' , contradicting the definition of \tilde{V} . Here, we assume that $n_0 > n_0(\gamma/4, H')$. Note, that for each $v \in V$ there could be several copies of H' with v playing a role of v' . We choose one such copy H'_v arbitrarily. Let $V \subseteq \tilde{V}$ be a set of vertices whose degree in G is at least half the average degree of G , i.e. for each $y \in V$,

$$d_G(y) \geq \frac{\gamma}{2} \frac{\|L_j\|}{|V_j|}.$$

The total number of edges of G not incident to V is at most $\frac{1}{4}\gamma\|L_j\| + \frac{1}{2}\|G\| = \frac{3}{4}\gamma\|L_j\|$. Thus $\|G[V, V_{j-1}]\| \geq \frac{\gamma}{4}\|L_j\|$. Since all vertices from V_j in L_j have the same degree,

$$|V| \geq \frac{\gamma}{4}|V_j|.$$

Let for each $v \in V$, $A_v \subseteq V_j$ and $B_v \subseteq V_{j-1}$ be sets of vertices such that $V(H'_v) = A_v \cup B_v \cup \{v\}$, $v \notin A_v$. Randomly color each vertex in V with red or blue independently with equal probability, color the vertices in $V_j - V$ blue. We say that a vertex $v \in V$ is *good* if v is red and each vertex in A_v is blue. Then the expected number of good v 's is at least $|V|2^{-t}$, where $|A_v| = t - 1$. Thus there is a set $V' \subseteq V$, corresponding to a set of good v 's, with $|V'| \geq |V|2^{-t}$, such that for each $v \in V'$, $A_v \cap V' = \emptyset$.

We see that

$$\|G[V', V_{j-1}]\| \geq \frac{\gamma}{2} \frac{\|L_j\|}{|V_j|} |V'| \geq \frac{\gamma}{2} \frac{\|L_j\|}{|V_j|} \frac{|V|}{2^t} \geq \gamma^2 2^{-t-3} \|L_j\|.$$

Consider the graph F defined above. By Lemma 3 F has a partite representation. Then by Lemma 4 there is a copy F^* of F in $G[V' \cup V_{j-1}]$ with a vertex $v \in V$ corresponding to v' . Here, we assume that $n_0 > n_1(\gamma^2 2^{-t-3}, F)$. Let F_i^* be a respective copy of F_i , for each $i \in [q]$. By construction of V' , there is a copy H^* of H' in G with all vertices except for v not in V' . Let B_v be the set of vertices of H^* in V_{j-1} . We see that $V(F^*) \cap V(H^*) \cap V_j = \{v\}$. Since $q > |V(H')| > |B_v|$, there is $i \in [q]$, such that $V(F_i^*) \cap B_v \cap V_{j-1} = \emptyset$. Then $H^* \cup F_i^*$ is a copy of H in G .

The case when v' is not an odd vertex is treated similarly by first finding many copies of H' with a vertex corresponding to v' in V_{j-1} . \square

Proof of Theorem 1. Let G' be a subgraph of Q_n such that $\|G'\| = 2\gamma\|Q_n\|$, for some constant $\gamma > 0$ and sufficiently large n . By a standard argument, see for example Lemma 1 [4],

$$\sum_{i:|i-n/2|>n^{2/3}} \binom{n}{i} = o(2^n).$$

Since the degree of each vertex in Q_n is n , the total number of edges in G' , incident to vertices in V_i 's, for $i < n/2 - n^{2/3}$ or $i > n/2 + n^{2/3}$ is $o(n2^n) = o(\|Q_n\|) = o(\|G'\|)$. Then there is $j \in \{n/2 - n^{2/3}, n/2 + n^{2/3}\}$ such that L_j contains at least $\gamma\|L_j\|$ edges of G' . Lemma 5 applied to $G = G' \cap L_j$ concludes the proof. \square

Proof of Theorem 2. Let $H = H(q)$ be a union of two copies Θ_1 and Θ_2 of $\Theta(q)$ sharing exactly one vertex that is a main pole of Θ_2 and a subdivision vertex of Θ_1 .

First we need to check that $H(q)$ is cubical. Marquardt [15] gave an explicit embedding of $H(3)$ in a hypercube. One can also use a characterisation by Havel and Moravek [14], who proved that a graph is cubical if and only if it has a nice edge-coloring. Here, an edge-coloring is *nice* if any cycle uses each color an even number of times and each non-trivial path uses some color an odd number of times. To construct a nice coloring of H , one can use nice colorings of Θ_1 and Θ_2 using disjoint sets of colors. These colorings were given in [14] and we show one in Figure 1.

Next, we shall show that $H(q)$ has no partite representation. A Hamming distance between two binary vectors a and b , denoted $d_H(a, b)$ is the number of positions where the vectors differ. In [2] it is shown that if $q \geq 3$ and a, a' are the main poles of a copy of $\Theta(q)$ embedded in a layer of a hypercube, then $d_H(a, a') = 2$. Let the main poles of Θ_1 and Θ_2 be denoted v'_i, v''_i , respectively, $i = 1, 2$. Assume that H has a k -partite representation for some k , i.e., $V(H) \subseteq \binom{[n]}{k} \cup \binom{[n]}{k-1}$, for some n and $V(H) \cap \binom{[n]}{k}$ corresponds to the edges of a k -partite hypergraph, denote it by \mathcal{H} . We have that $d_H(v'_i, v''_i) = 2$, $i = 1, 2$. Since a main pole of Θ_1 and a main pole of Θ_2 are adjacent, the main poles of Θ_1 are in one vertex layer and the main poles of Θ_2 are in another vertex layer, without loss of generality, $v'_1, v''_1 \in \binom{[n]}{k}$ and $v'_2, v''_2 \in \binom{[n]}{k-1}$. Assume further that v'_1 and v''_1 equal to 10 and 01 in the first two positions of their binary representation and coincide on all other positions. Consider neighbours of v_1 in Θ_1 . At least one of these neighbours, say w differs from v'_1 in a position that is not one of the first two ones, say in the third position. Then, restricted to the first three position v'_1 is 101, w is 100, and v''_1 is 011. Thus, the remaining two vertices on the v_1, v'_1 path, that contains w , must be equal to 110 and 010 in these three position. All vertices of this leg are the same in positions $4, \dots, n$, say they are equal to 1 on a set of positions $A \subseteq \{4, \dots, n\}$, $|A| = k - 2$. Thus, we have that \mathcal{H}' contains hyperedges $\{1, 2\} \cup A$, $\{2, 3\} \cup A$, and $\{1, 3\} \cup A$. If \mathcal{H} were to be k -partite, 1, 2, 3 and each element of A would belong to distinct parts, i.e., to $k + 1$ parts, a contradiction. This shows that $H(q)$ has no partite representation.

Finally, we shall show that $H(q)$ has a zero Turán density in a hypercube. Using Theorem 1 it is sufficient to show that $\Theta(q)$ has partite representation. Note that $\Theta(q) = K_{2,q}(1)$, a subdivision of $K_{2,q}$. We claim that $\mathcal{H} = K_{2,q}$ gives a 2-partite representation of $\Theta(q)$. Indeed, let $n = q + 2$ and let \mathcal{H} be a hypergraph on a vertex set $[n]$ with the edge set $\{ij : i \in [2], j \in [n] \setminus [2]\}$. So, \mathcal{H} is a bipartite graph, i.e., 2-partite 2-uniform hypergraph. Let the edges of \mathcal{H} correspond to subdivision vertices of $K_{2,q}(1)$. Let vertices 1 and 2 correspond to the main poles of $\Theta(q)$ and vertices $2, \dots, q$ correspond to other vertices. This concludes the proof of Theorem 2. \square

Conclusions We showed that there are cubical graphs that have no partite representation and have zero Turán density in a hypercube. On the other hand we proved that any graph whose every block has a partite representation has a zero Turán density in a hypercube. This leads to a followup question:

Open question Is it true that in each cubical graph that has a zero Turán density in a hypercube each block has a partite representation?

Acknowledgements The research was supported in part by a DFG grant FKZ AX 93/2-1.

References

- [1] N. Alon, A. Krech and T. Szabó. Turán’s theorem in the hypercube. *SIAM Journal on Discrete Mathematics*, 21(1) (2007), 66–72.
- [2] M. Axenovich, R. Martin, C. Winter. On graphs embeddable in a layer of a hypercube and their extremal numbers. ArXiv preprint 2023: arXiv:2303.15529
- [3] M. Axenovich. Extremal numbers for cycles in a hypercube. To appear in *Discrete Applied Mathematics*, ArXiv preprint 2022: arXiv:2211.12842
- [4] M. Axenovich, J. Manske, R. Martin, Extremal functions for Q_2 in a Boolean lattice, *Order*, Online First, 14 March 2011.
- [5] R. Baber. Turán densities of hypercubes. Arxiv preprint (2012) arXiv:1201.3587.
- [6] J. Balogh, P. Hu, B. Lidický and H. Liu. Upper bounds on the size of 4- and 6-cycle-free subgraphs of the hypercube. *European Journal of Combinatorics* 35 (2014), 75–85.
- [7] P. Brass, H. Harborth and H. Nienborg. On the maximum number of edges in a C_4 -free subgraph of Q_n . *Journal of Graph Theory* 19, (1995) 17–23.
- [8] F. Chung. Subgraphs of a hypercube containing no small even cycles. *Journal Graph Theory* 16 (1992), 273–286.
- [9] M. Conder. Hexagon-free subgraphs of hypercubes. *Journal of Graph Theory* 17 (4) (1993), 477–479.
- [10] D. Conlon. An extremal theorem in the hypercube. *Electronic Journal of Combinatorics* 17 (2010), no. R111.
- [11] P. Erdős. On extremal problems of graphs and generalized graphs. *Israel J. Math.* 2 (1964) 183–190.
- [12] Z. Füredi and L. Özkahya. On even-cycle-free subgraphs of the hypercube. *Electronic Notes in Discrete Mathematics* 34 (2009), 515–517.
- [13] Z. Füredi and L. Özkahya. On even-cycle-free subgraphs of the hypercube. *Journal of Combinatorial Theory, Series A*, 118 (2011), 1816–1819.
- [14] I. Havel and J. Morávek. B-valuations of graphs. *Czechoslovak Mathematical Journal*, Vol. 22 (1972), No. 2, 338–351.
- [15] F. Marquardt. On the characterization of graphs with zero Turán density in the hypercube. Master thesis, Karlsruhe Institute of Technology, 2022.
- [16] D. Offner. Some Turán type results on the hypercube. *Discrete Mathematics*, 309(9) (2009), 2905–2912.
- [17] A. Thomason and P. Wagner. Bounding the size of square-free subgraphs of the hypercube. *Discrete Math.* 309 (2009), 1730–1735.
- [18] I. Tomon. Robust (rainbow) subdivisions and simplicial cycles. ArXiv preprint 2022: arXiv:2201.12309.