

NOTES FROM THE WORKSHOP ON APPROXIMATE GROUPS AND APERIODIC ORDER

MATTIAS BYLÉHN

Based on joint work with Michael Björklund

1. HEAT KERNEL HYPERUNIFORMITY

Let Λ be an invariant (locally square-summable) point process on hyperbolic space

$$\mathbb{H}^n = \mathrm{SO}^+(1, n)/\mathrm{SO}(n)$$

and let σ_Λ denote the associated (centered) diffraction measure on $(0, +\infty) \cup i[0, \rho) \subset \mathbb{C}$, where $\rho = (n-1)/2$. Recall from the talk that σ_Λ is the unique positive Radon measure on $(0, +\infty) \cup i[0, \rho)$ satisfying

$$\mathrm{Var}\left(\sum_{p \in \Lambda} f(p)\right) = \int_0^\infty |\widehat{f}(\lambda)|^2 d\sigma_\Lambda^{(p)}(\lambda) + \int_0^\rho |\widehat{f}(i\lambda)|^2 d\sigma_\Lambda^{(c)}(\lambda)$$

for all compactly supported continuous functions f on \mathbb{H}^n , where $\sigma_\Lambda^{(p)}, \sigma_\Lambda^{(c)}$ are the restrictions of σ_Λ to $(0, +\infty)$ and $i[0, \rho)$ respectively.

Assumption: We will assume throughout this note that $\sigma_\Lambda^{(c)} = 0$, saving us from any "super-Poissonian" density fluctuations of the process.

Goal: We want show that such a point process Λ is spectrally hyperuniform, i.e.

$$\sigma_\Lambda([0, \varepsilon]) = o(\varepsilon^3),$$

if and only if

$$\limsup_{t \rightarrow +\infty} t^{3/2} e^{2\rho^2 t} \mathrm{Var}\left(\sum_{p \in \Lambda} h_t(p)\right) = 0, \quad (1.1)$$

where

$$h_t(x) = \int_0^\infty e^{-t(\rho^2 + \lambda^2)} \phi_\lambda(d(x, o)) \frac{d\lambda}{|c_n(\lambda)|^2}$$

is the heat kernel on \mathbb{H}^n . If Λ satisfies equation (1.1), then we say that it is *heat kernel hyperuniform*.

1.1. Spectral hyperuniformity implies heat kernel hyperuniformity

Assume that Λ is spectrally hyperuniform. We will first rewrite the variance of the periodized heat kernel into a suitable form and then formulate Lemma 1.1, where the heart of the proof lies.

Consider the function $f_t = e^{\rho^2 t} h_t$ on \mathbb{H}^n , so that

$$\widehat{f}_t(\lambda) = e^{\rho^2 t} \widehat{h}_t(\lambda) = e^{\rho^2 t} e^{-t(\rho^2 + \lambda^2)} = e^{-t\lambda^2} =: \psi(\sqrt{t}\lambda).$$

Using the diffraction measure σ_Λ , we rewrite the heat kernel variance as

$$t^{3/2} e^{2\rho^2 t} \mathrm{Var}\left(\sum_{p \in \Lambda} h_t(p)\right) = t^{3/2} \mathrm{Var}\left(\sum_{p \in \Lambda} f_t(p)\right) = t^{3/2} \int_0^\infty \psi(\sqrt{t}\lambda)^2 d\sigma_\Lambda(\lambda).$$

Letting $s = \sqrt{t}$, we have to show that

$$s^3 \int_0^\infty \psi(s\lambda)^2 d\sigma_\Lambda(\lambda) \longrightarrow 0, \quad s \rightarrow +\infty.$$

Set

$$\varphi(r) = -2\psi'(r)\psi(r) = 4r e^{-2r^2}$$

so that

$$\psi(s\lambda)^2 = s \int_\lambda^\infty \varphi(sr) dr.$$

By a standard change of variables,

$$\begin{aligned} \int_0^\infty \psi(s\lambda)^2 d\sigma_\Lambda(\lambda) &= s \int_0^\infty \left(\int_\lambda^\infty \varphi(sr) dr \right) d\sigma_\Lambda(\lambda) = s \int_0^\infty \varphi(sr) \left(\int_0^t d\sigma_\Lambda(\lambda) \right) dr \\ &= s \int_0^\infty \varphi(sr) \sigma_\Lambda([0, r]) dr. \end{aligned}$$

This means that it is enough to show that

$$s^4 \int_0^\infty \varphi(sr) \sigma_\Lambda([0, r]) dr \longrightarrow 0, \quad s \rightarrow +\infty.$$

The core of the proof lies in the following Lemma.

Lemma 1.1 (Björklund). *Suppose that there are constants $\alpha, \beta > 0$, $\gamma \in (0, 1)$, $A, B, C \geq 0$ and Borel functions $\omega, \varphi : [0, +\infty) \rightarrow [0, \infty)$ satisfying*

(i) ρ is increasing and such that

- for every $\varepsilon > 0$ there is an $r_\varepsilon \in (0, 1)$ with $r_\varepsilon \rightarrow 0$ such that $\omega(r) \leq A\varepsilon r^\alpha$ for all $r \in [0, r_\varepsilon^{1-\gamma}]$,
- there is a constant $M > 0$ such that $\omega(r) \leq Br^\beta$ for all $r \geq M$.

(ii) φ is such that

- (1) $\int_0^\infty r^\alpha \varphi(r) dr < +\infty$,
- (2) $R^{\alpha/\gamma} \int_R^\infty \varphi(r) dr \rightarrow 0$ as $R \rightarrow +\infty$,
- (3) $R^{\alpha-\beta} \int_R^\infty r^\beta \varphi(r) dr \rightarrow 0$ as $R \rightarrow +\infty$.

Then

$$r_\varepsilon^{-(\alpha+1)} \int_0^\infty \varphi(r/r_\varepsilon) \omega(r) dr \longrightarrow 0, \quad \varepsilon \rightarrow 0^+.$$

With this Lemma in mind, we set $\varphi(r) = 4re^{-2r^2}$ and $\omega(r) = \sigma_\Lambda([0, r])$. By the assumed spectral hyperuniformity, there is a constant $A \geq 0$ and a sequence $r_\varepsilon \rightarrow 0$ such that

$$\omega(r) \leq A\varepsilon r^3, \quad r \in [0, r_\varepsilon^{1/2}],$$

so we can take $\alpha = 3$, $\gamma = 1/2$. Finding constants B, β as in the Lemma above requires some more work.

Lemma 1.2. *Let Λ be an invariant point process on \mathbb{H}^n with diffraction σ_Λ . Then $\sigma_\Lambda([0, R]) \ll R^n$ for sufficiently large $R > 0$.*

Remark 1.3. When $\omega(r) = \sigma([0, r])$ is the diffraction measure of a point process, Lemma 1.1 is applicable for all rank one spaces. In particular, the crucial constants for Euclidean and hyperbolic space are

G/K	\mathbb{R}^n	\mathbb{H}^n
α	n	3
β	n	n

Thus for hyperbolic space there is a $B > 0$ and an $M > 0$ such that $\rho(r) \leq Br^n$ for all $r \geq M$, so take $\beta = n$.

Modulo the proof of Lemma 1.1, it suffices to verify points (1)-(3) for the function $\varphi(r) = 4re^{-2r^2}$.

(1) we have

$$4 \int_0^\infty r^4 e^{-2r^2} dr < +\infty.$$

(2)

$$R^6 \int_R^\infty 4re^{-2r^2} dr = 2R^6 e^{-2R^2} \rightarrow 0, \quad R \rightarrow +\infty.$$

(3)

$$R^{3-n} \int_R^\infty r^n e^{-2r^2} dr = R^4 \int_1^\infty r^n e^{-2R^2 r^2} dr \rightarrow 0, \quad R \rightarrow +\infty$$

by dominated convergence.

Letting $s_\varepsilon = r_\varepsilon^{-1}$, then Lemma 1.1 tells us that

$$s_\varepsilon^4 \int_0^\infty \varphi(s_\varepsilon r) \sigma_\Lambda([0, r]) dr \rightarrow 0, \quad \varepsilon \rightarrow 0^+,$$

which is what we wanted to show.

Proof of Lemma 1.1. Let $I_\varepsilon = \int_0^\infty \varphi(r/r_\varepsilon) \rho(r) dr$. Then we use the assumed monotone growth and estimates on ω to find that

$$I_\varepsilon \leq \int_0^{r_\varepsilon^{1-\gamma}} \varphi(r/r_\varepsilon) A \varepsilon r^\alpha dr + \omega(M) \int_{r_\varepsilon^{1-\gamma}}^M \varphi(r/r_\varepsilon) dr + \int_M^\infty \varphi(r/r_\varepsilon) B r^\beta dr =: I_1 + I_2 + I_3.$$

We need to show that $I_j = o(r_\varepsilon^{\alpha+1})$ for $j = 1, 2, 3$ as $\varepsilon \rightarrow 0^+$.

$$\begin{aligned} I_1 &\leq A \varepsilon r_\varepsilon^{\alpha+1} \int_0^\infty \varphi(r) r^\alpha dr \\ I_2 &\leq \varphi(M) r_\varepsilon \int_{r_\varepsilon^{-\gamma}}^\infty \varphi(r) dr \\ I_3 &= B r_\varepsilon^{\beta+1} \int_{M/r_\varepsilon}^\infty \varphi(r) r^\beta dr. \end{aligned}$$

Dividing by $r_\varepsilon^{\alpha+1}$, we get that

$$\frac{I_\varepsilon}{r_\varepsilon^{\alpha+1}} = \frac{I_1 + I_2 + I_3}{r_\varepsilon^{\alpha+1}} \ll \underbrace{\varepsilon + \omega(M) (r_\varepsilon^{-\gamma})^{\alpha/\gamma} \int_{r_\varepsilon^{-\gamma}}^\infty \varphi(r) dr}_{\rightarrow 0} + \underbrace{r_\varepsilon^{-(\alpha-\beta)} \int_{M/r_\varepsilon}^\infty \varphi(r) r^\beta dr}_{\rightarrow 0}.$$

■

1.2. Heat kernel hyperuniformity implies spectral hyperuniformity

Assume that equation (1.1) holds. We will show that $\sigma_\Lambda([0, \varepsilon]) = o(\varepsilon^3)$ as $\varepsilon \rightarrow 0^+$.

This is fortunately quite straightforward: Let $\delta_t > 0$ and bound canonically from above,

$$\text{Var}\left(\sum_{p \in \Lambda} h_t(p)\right) = \int_0^\infty |\widehat{h}_t(\lambda)|^2 d\sigma_\Lambda(\lambda) \geq \int_0^{\delta_t} |\widehat{h}_t(\lambda)|^2 d\sigma_\Lambda(\lambda) = e^{-2\rho^2 t} \int_0^{\delta_t} e^{-2t\lambda^2} d\sigma_\Lambda(\lambda).$$

Assuming heat kernel hyperuniformity, there is for every $\varepsilon > 0$ a $t_\varepsilon > 0$ such that

$$t^{3/2} \int_0^{\delta_t} e^{-2t\lambda^2} d\sigma_\Lambda(\lambda) < \varepsilon, \quad \forall t \geq t_\varepsilon.$$

In particular,

$$t^{3/2} e^{-2t\delta_t^2} \sigma_\Lambda([0, \delta_t]) < \varepsilon, \quad \forall t \geq t_\varepsilon,$$

so the question boils down to the following: If we can find $\delta_t \rightarrow 0$ as $t \rightarrow +\infty$ such that

$$\delta_t^{-3} \ll t^{3/2} e^{-2t\delta_t^2}, \quad \forall t \geq t_\varepsilon, \tag{1.2}$$

then

$$\frac{\sigma_\Lambda([0, \delta_t])}{\delta_t^3} \ll t^{3/2} e^{-2t\delta_t^2} \sigma_\Lambda([0, \delta_t]) < \varepsilon, \quad \forall t \geq t_\varepsilon,$$

as desired.

Finding a solution δ_t to equation (1.2) is equivalent to $2t\delta_t^2 - 3 \log(\delta_t) \leq \frac{3c}{2} \log(t)$ for some $c > 0$ for all sufficiently large t . As an example, by setting $c = 2$ we get $2t\delta_t^2 \leq 3 \log(t\delta_t)$, for which $\delta_t = t^{-1/2}$ is an example of a solution whenever $t \geq e^{4/3}$.