

---

# BOLLETTINO UNIONE MATEMATICA ITALIANA

---

MICHAEL STRUWE, GABRIELLA TARANTELLA

## On multivortex solutions in Chern-Simons gauge theory

*Bollettino dell'Unione Matematica Italiana, Serie 8, Vol. 1-B (1998),  
n.1, p. 109–121.*

Unione Matematica Italiana

[http://www.bdim.eu/item?id=BUMI\\_1998\\_8\\_1B\\_1\\_109\\_0](http://www.bdim.eu/item?id=BUMI_1998_8_1B_1_109_0)

L'utilizzo e la stampa di questo documento digitale è consentito liberamente per motivi di ricerca e studio. Non è consentito l'utilizzo dello stesso per motivi commerciali. Tutte le copie di questo documento devono riportare questo avvertimento.

---

*Articolo digitalizzato nel quadro del programma  
bdim (Biblioteca Digitale Italiana di Matematica)  
SIMAI & UMI*

<http://www.bdim.eu/>



## On Multivortex Solutions in Chern-Simons Gauge Theory.

MICHAEL STRUWE - GABRIELLA TARANTELLO

**Sunto.** – *Motivati dall'analisi asintotica dei vortici nella teoria di Chern-Simons-Higgs, si studia l'equazione*

$$-\Delta u = \lambda \left( \frac{e^u}{\int_{\Omega} e^u dx} - \frac{1}{|\Omega|} \right), \quad u \in H^1(\Omega)$$

dove  $\Omega = \mathbb{R}^2 / \mathbb{Z}^2$  è il toro piatto bidimensionale. In contrasto con l'analogo problema di Dirichlet, si dimostra che per  $\lambda \in ]8\pi, 4\pi^2[$  l'equazione ammette una soluzione non banale. Tale soluzione cattura il carattere bidimensionale dell'equazione, nel senso che, per tali valori di  $\lambda$ , l'equazione non può ammettere soluzioni (periodiche) non banali dipendenti da una sola variabile (vedi [10]).

### 1. – Introduction.

In  $(2+1)$ -dimensional Chern-Simons gauge theory, a particular role is played by the corresponding condensate (or multivortex) solutions which are believed relevant in several aspects of theoretical physics.

Although the presence of multivortices has been predicted experimentally, still in the general framework of Chern-Simons theory, it is very difficult to obtain them analytically. Thus, a special effort has been devoted to derive specific models for which more convenient selfdual equations would hold for the corresponding energy-minimizing multivortices. See the recent monograph [4].

Through an approach of Taubes [15], the process of solving these selfdual equations is reduced to solving suitable elliptic equations for the logarithmic values of the particle density. The elliptic equations so derived are of Liouville-type. It is necessary to solve them on the 2-dimensional torus in order to obtain the desired condensate solution subject to 't Hooft periodic boundary conditions.

Here we consider a particular class of these equations which were derived in [14]. More precisely, [14] is concerned with a selfdual model introduced in [5] and [6]; it establishes the existence of a new class of condensate solutions which are absent in the classical vortex theory.

When the vortex number  $N=1$  and the Chern-Simons coupling constant tends to zero, in [14] it is shown that the asymptotic behavior of the new type of

condensates can be described in terms of solutions of the limiting equation

$$(1) \quad -\Delta u = \lambda \left( \frac{e^{w_0+u}}{\int_{\Omega} e^{w_0+u}} - \frac{1}{|\Omega|} \right) \quad \text{on } \Omega$$

with  $\lambda = 4\pi$ ,  $\Omega$  the 2-dimensional torus, and  $w_0$  an assigned function.

Notice that for  $\lambda \in ]0, 8\pi[$  existence for (1) is an easy consequence of the Moser-Trudinger inequality [9]. To extend this argument to condensate solutions with vortex number  $N \geq 2$ , it is necessary to insure the existence of solutions for (1) when  $\lambda \geq 8\pi$ . This is exactly the task we have taken up here. We treat the case  $w_0 = 0$  and show that (1) admits a nonconstant solution for every  $\lambda$  in the range  $8\pi < \lambda < 4\pi^2$ .

By a result of Ricciardi-Tarantello [10], we can also guarantee that these solutions are «truly» two-dimensional in the sense that they cannot reduce to a (periodic) function of one variable. Incidentally, let us also point out that the analogous problem (with  $w_0 = 0$ ) subject to Dirichlet boundary conditions

$$(2) \quad \begin{cases} -\Delta u = \lambda \frac{e^u}{\int_{\Omega} e^u} & \text{on } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where  $\Omega \subset \mathbb{R}^2$  is a bounded domain, plays an important role in the context of statistical mechanics of point vortices in the mean field limit where (2) is referred to as the mean field equation; see [2], [3], [7].

Since for  $0 < \lambda < 8\pi$  and  $\Omega$  simply connected (2) is known to admit a unique solution (see [13]), by analogy one would be tempted to conjecture that problem (1) with  $w_0 = 0$  and  $0 < \lambda < 8\pi$  also admits only the trivial solution  $u = 0$ . We can establish this result only for  $\lambda$  small (see section 5) but we are not certain about its validity in the whole range  $]0, 8\pi[$ . In fact, our result shows that there is an important difference between problems (1) (with  $w_0 = 0$ ) and (2), as problem (2) admits no solutions for  $\lambda \geq 8\pi$ , if  $\Omega$  is a ball.

*Acknowledgements.* The authors wish to thank Y. Y. Li and S. Müller for useful discussions.

## 2. – Main result.

Let  $\Omega$  be the 2-dimensional torus, with fundamental cell domain:  $[-1/2, 1/2] \times [-1/2, 1/2]$ . Consider the problem

$$(3) \quad -\Delta u = \lambda \left( \frac{e^u}{\int_{\Omega} e^u dx} - 1 \right) \quad \text{on } \Omega,$$

or, equivalently, solutions of (3) on  $\mathbb{R}^2$  of period 1 in each variable. For fixed  $\lambda$ , we refer to equation (3) as  $(3)_\lambda$ . Shifting a solution  $u$  of  $(3)_\lambda$  by a constant, we again obtain a solution. We normalize solutions by requiring  $\int_{\Omega} u \, dx = 0$ .

Notice that  $u = 0$  is always a solution of  $(3)_\lambda$ ; here we seek nontrivial solutions.

Let  $E = \{u \in H^1(\Omega); \int_{\Omega} u \, dx = 0\}$  with norm  $\|u\| = \int_{\Omega} |\nabla u|^2 \, dx$ . Then (weak) solutions of  $(3)_\lambda$  correspond to critical points of the analytic functional

$$I_\lambda(u) = \frac{1}{2} \|u\|^2 - \lambda \ln \left( \int_{\Omega} e^u \, dx \right) \quad \text{on } E.$$

REMARK 2.1. – By Jensen’s inequality we have  $\int_{\Omega} e^u \, dx \geq e^{\int_{\Omega} u \, dx} = 1$  for all  $u \in E$ ; in particular, the map  $\lambda \rightarrow I_\lambda(u)$  is monotone decreasing for any  $u \in E$ .

REMARK 2.2. – By Trudinger-Moser’s inequality [9], it is easy to check that,  $I_\lambda$  is bounded from below, coercive, and weakly lower semicontinuous if  $\lambda < 8\pi$ . So  $I_\lambda$  achieves its infimum, which, however, could correspond to the trivial solution  $u = 0$ .

On the other hand, we shall see that for  $\lambda > 8\pi$  the functional  $I_\lambda$  is unbounded from below, while the trivial solution  $u \equiv 0$  remains a strict local minimum for  $\lambda < 4\pi^2$ . Thus, for  $8\pi < \lambda < 4\pi^2$  the functional  $I_\lambda$  exhibits a mountain-pass structure and we expect the existence of non-trivial critical points of  $I_\lambda$  for  $\lambda$  in this range. This, in fact, is our main result.

THEOREM 2.1. – *For every  $\lambda \in ]8\pi, 4\pi^2[$  there exists a non-trivial solution  $u_\lambda$  of  $(3)_\lambda$  satisfying  $I_\lambda(u_\lambda) \geq (1 - \lambda/4\pi^2) c_0$  for some constant  $c_0 > 0$  independent of  $\lambda$ .*

The solutions  $u_\lambda$  will be obtained by a variational method using, in particular, the strategy of obtaining a priori bounds on Palais-Smale sequences by parameter variation, as introduced in [11], [12]. We expect these solutions to form a continuous «branch», bifurcating from the trivial branch  $u \equiv 0$  at  $\lambda = 4\pi^2$  and asymptotic to the line  $\lambda = 8\pi$ . However, at this stage we cannot rigorously prove that this is the case. Moreover, we do not know if non-trivial solutions also exist for  $\lambda \leq 8\pi$ , in particular, for  $\lambda = 8\pi$ , but some analytical evidence seems to suggest that they do.

We also point out that the solution  $u_\lambda$  cannot reduce to a (periodic) function of one variable. In fact, for the corresponding one-dimensional problem:

$$(4) \quad \ddot{u} + \lambda \left( \frac{e^u}{\int_{-1/2}^{1/2} e^u \, dx} - 1 \right) = 0$$

a recent result of Ricciardi-Tarantello [10] asserts that (4) admits a nonconstant solution of period  $T = 1$  if and only if  $\lambda > 4\pi^2$ . Thus, Theorem 2.1 captures, in an

essential way, the two-dimensional nature of problem  $(3)_\lambda$  and this justifies the special role played by the value  $\lambda = 4\pi^2$ .

### 3. – Existence of solutions for almost every $\lambda$ .

In a first step we show that nontrivial solutions to  $(3)_\lambda$  exist for almost every  $\lambda \in ]8\pi, 4\pi^2[$ .

LEMMA 3.1. – *If  $\lambda < 4\pi^2$ , then  $u = 0$  is a strict local minimum for  $I_\lambda$ .*

PROOF. –  $I_\lambda$  is smooth. Thus it suffices to observe that the second variation of  $I_\lambda$  at  $u = 0$  in direction  $v \in E$  can be estimated

$$I_\lambda''(0)(v, v) = \|v\|^2 - \lambda \int_{\Omega} v^2 dx \geq \left(1 - \frac{\lambda}{4\pi^2}\right) \|v\|^2. \quad \blacksquare$$

For  $\varepsilon > 0$  and  $x \in \Omega$  let

$$v_\varepsilon(x) = \ln \left( \frac{\varepsilon^2}{(\varepsilon^2 + \pi|x|^2)^2} \right),$$

extended periodically, and let  $u_\varepsilon = v_\varepsilon - \int_{\Omega} v_\varepsilon dx \in E$ .

LEMMA 3.2. –  $I_\lambda(u_\varepsilon) = 2(8\pi - \lambda) \ln(1/\varepsilon) + O(1)$ , where  $|O(1)| \leq C$  as  $\varepsilon \rightarrow 0$ .

PROOF. – We estimate

$$|\nabla u_\varepsilon|^2 = 4 |\nabla \ln(\varepsilon^2 + \pi|x|^2)|^2 = \frac{16\pi^2 |x|^2}{(\varepsilon^2 + \pi|x|^2)^2}.$$

Substituting  $y = x/\varepsilon$ , we obtain

$$\|u_\varepsilon\|^2 = 16\pi^2 \int_{\Omega_\varepsilon} \frac{|y|^2}{(1 + \pi|y|^2)^2} dy,$$

where  $\Omega_\varepsilon = \{y; \varepsilon y \in \Omega\}$ . Introducing polar coordinates around 0, the latter equals

$$\|u_\varepsilon\|^2 = 32\pi^3 \int_0^{\varepsilon^{-1}} \frac{r^3 dr}{(1 + \pi r^2)^2} + O(1) = 32\pi \ln \frac{1}{\varepsilon} + O(1),$$

where  $|O(1)| \leq C$  for  $\varepsilon \rightarrow 0$ .

On the other hand, we have

$$\ln \left( \int_{\Omega} e^{u_\varepsilon} dx \right) = \ln \left( \int_{\Omega} e^{v_\varepsilon} dx \right) - \int_{\Omega} v_\varepsilon dx,$$

and

$$\int_{\Omega} e^{v_{\varepsilon}} dx = \int_{\Omega} \frac{\varepsilon^2 dx}{(\varepsilon^2 + \pi|x|^2)^2} = \int_{\Omega_{\varepsilon}} \frac{dy}{(1 + \pi|y|^2)^2} = O(1),$$

while

$$\int_{\Omega} v_{\varepsilon} dx = \int_{\Omega} \ln \left( \frac{\varepsilon^2}{(\varepsilon^2 + \pi|x|^2)^2} \right) dx = 2 \ln \varepsilon - 2 \int_{\Omega} \ln (\varepsilon^2 + \pi|x|^2) dx = 2 \ln \varepsilon + O(1).$$

Thus, we obtain, with  $|O(1)| \leq C$  as  $\varepsilon \rightarrow 0$ , the estimate

$$I_{\lambda}(u_{\varepsilon}) = \frac{1}{2} \|u_{\varepsilon}\|^2 - \lambda \ln \left( \int_{\Omega} e^{u_{\varepsilon}} dx \right) = (16\pi - 2\lambda) \ln \frac{1}{\varepsilon} + O(1),$$

as desired. ■

REMARK 3.1. – Note, in particular, that  $\|u_{\varepsilon}\| \rightarrow \infty$  as  $\varepsilon \rightarrow 0$ .

Fix  $\lambda \in ]8\pi, 4\pi^2[$ . By Lemma 3.2 there exists  $\varepsilon_0 = \varepsilon_0(\lambda) > 0$  sufficiently small such that for  $u_0 = u_{\varepsilon_0}$  we have

$$I_{\lambda}(u_0) < 0 \quad \text{and} \quad \|u_0\| \geq 1.$$

Hence also for any  $\mu \geq \lambda$  we have  $I_{\mu}(u_0) \leq I_{\lambda}(u_0) < 0$ .

Define

$$P = \{ \gamma : [0, 1] \rightarrow E; \gamma \text{ is continuous } \gamma(0) = 0, \gamma(1) = u_0 \}$$

and for  $\mu \geq \lambda$  let

$$c_{\mu} = \inf_{\gamma \in P} \max_{t \in [0, 1]} I_{\mu}(\gamma(t)).$$

In view of Remark 2.1, the map  $\mu \rightarrow c_{\mu}$ ,  $\mu \geq \lambda$  is monotone decreasing, hence differentiable at almost all values  $\mu \in ]\lambda, 4\pi^2[$ .

In addition, by Lemma 3.1, there exists a constant  $c_0 > 0$  (independent of  $\lambda$ ) such that

$$c_{\mu} \geq \left( 1 - \frac{\mu}{4\pi^2} \right) c_0.$$

LEMMA 3.3. – Suppose the map:  $\mu \rightarrow c_{\mu}$  is differentiable at  $\mu > \lambda$ . Then  $c_{\mu}$  defines a critical value for  $I_{\mu}$ . In particular, problem (3) <sub>$\mu$</sub>  admits a nontrivial solution for almost every  $\mu \in ]\lambda, 4\pi^2[$ .

PROOF. – Let  $\mu$  be a point of differentiability of  $c_{\mu}$ . Consider a monotone decreasing sequence  $(\mu_n)$  such that  $\mu_n \rightarrow \mu (n \rightarrow \infty)$ . For  $n \in \mathbb{N}$  and any path  $\gamma_n \in P$

such that

$$(5) \quad \max_{t \in [0, 1]} I_\mu(\gamma_n(t)) \leq c_\mu + (\mu_n - \mu)$$

consider any point  $u = \gamma_n(t_n)$  such that  $I_{\mu_n}(u) \geq c_{\mu_n} - 2(\mu_n - \mu)$ .

Then, letting  $\alpha = -c'_\mu + 3$  and choosing  $n_0 \in \mathbb{N}$  sufficiently large, for  $n \geq n_0$  we have

$$(6) \quad c_\mu - \alpha(\mu_n - \mu) \leq c_{\mu_n} - 2(\mu_n - \mu) \leq I_{\mu_n}(u) \leq I_\mu(u) \leq \\ \leq \max_{0 \leq t \leq 1} I_\mu(\gamma_n(t)) \leq c_\mu + (\mu_n - \mu).$$

Note that  $n_0$  is independant of the choice of  $\gamma_n$ . In particular, (6) implies that

$$0 \leq \frac{I_\mu(u) - I_{\mu_n}(u)}{\mu_n - \mu} = \ln \left( \int_\Omega e^u dx \right) \leq \alpha + 1$$

and hence that

$$(7) \quad \|u\|^2 = 2I_\mu(u) + 2\mu \ln \left( \int_\Omega e^u dx \right) \leq 2c_\mu + 2(\mu_n - \mu) + 2\mu(\alpha + 1) \leq C_1$$

for any such point  $u = \gamma_n(t_n)$ , any  $n \geq n_0$ . To proceed, we need the following estimates.

LEMMA 3.4. – i) For any  $u, v \in E$ , any  $\mu \geq 0$  there holds

$$I_\mu(u + v) \leq I_\mu(u) + \langle I'_\mu(u), v \rangle + \frac{1}{2} \|v\|^2.$$

ii) For any  $C_1 \geq 0$  there exists a constant  $C$  such that for any  $\mu, v \in \mathbb{R}$  there holds

$$\|I'_\mu(u) - I'_\nu(u)\| \leq C|\mu - \nu|,$$

uniformly in  $u \in E$  with  $\|u\|^2 \leq C_1$ .

PROOF. – i) Expanding to second order, we have

$$I_\mu(u + v) - I_\mu(u) - \langle I'_\mu(u), v \rangle - \frac{1}{2} \|v\|^2 = \\ = -\mu \left\{ \ln \left( \frac{\int_\Omega e^{u+v} dx}{\int_\Omega e^u dx} \right) - \frac{\int_\Omega e^u v dx}{\int_\Omega e^u dx} \right\} = -\mu \int_0^1 \int_0^{s'} \frac{d^2 f}{ds^2}(s'') ds'' ds',$$

where  $f(s) = \ln \left( \int_\Omega e^{u+sv} dx / \int_\Omega e^u dx \right)$ .



Since by Schwarz' inequality

$$f''(s) = \frac{1}{\left(\int_{\Omega} e^{u+sv} dx\right)^2} \left\{ \int_{\Omega} e^{u+sv} v^2 dx \cdot \int_{\Omega} e^{u+sv} dx - \left(\int_{\Omega} e^{u+sv} v dx\right)^2 \right\} \geq 0,$$

the desired estimate follows.

ii) For any  $v \in E$  with  $\|v\| \leq 1$ , since  $\int_{\Omega} e^u dx \geq 1$ ,  $\|v\|_{L^2} \leq (1/2\pi)\|v\| \leq 1$ , we have

$$\begin{aligned} \langle I'_\mu(u), v \rangle - \langle I'_\nu(u), v \rangle &= (\nu - \mu) \frac{\int_{\Omega} e^u v dx}{\int_{\Omega} e^u dx} \leq |\mu - \nu| \left( \int_{\Omega} e^{2u} dx \cdot \int_{\Omega} v^2 dx \right)^{1/2} \leq \\ &\leq |\mu - \nu| \left( \int_{\Omega} e^{2u} dx \right)^{1/2} \leq e^{C_1/8\pi} |\mu - \nu| \left( \int_{\Omega} e^{4\pi(u^2/\|u\|^2)} dx \right)^{1/2}, \end{aligned}$$

where we used that

$$2|u| \leq 4\pi \frac{u^2}{\|u\|^2} + \frac{\|u\|^2}{4\pi} \leq 4\pi \frac{u^2}{\|u\|^2} + \frac{C_1}{4\pi}.$$

The claim now follows from the Trudinger-Moser inequality

$$\sup_{u \in E} \int_{\Omega} e^{4\pi(u^2/\|u\|^2)} dx < \infty;$$

see [9]. ■

Proceeding with the proof of Lemma 3.3, we can now construct a special (bounded) Palais-Smale sequence  $(u_n)$  for  $I_\mu$  at the energy level  $c_\mu$ .

LEMMA 3.5. - *There exists a sequence  $(u_n)$  in  $E$  such that  $\|u_n\|^2 \leq C_1$ ,  $I_\mu(u_n) \rightarrow c_\mu$  and  $I'_\mu(u_n) \rightarrow 0$  as  $n \rightarrow \infty$ .*

PROOF. - Otherwise, there exists  $\delta > 0$  such that  $\|I'_\mu(u)\| \geq 2\delta$  for all  $u \in E$  with  $\|u\|^2 \leq C_1$  and  $|I_\mu(u) - c_\mu| < 2\delta$ . We may assume that  $\alpha(\mu_n - \mu) < \delta$  for  $n \geq n_0$ .

Choose a function  $\varphi \in C^\infty(\mathbb{R})$  such that  $0 \leq \varphi \leq 1$ ,  $\varphi(s) = 1$  for  $s \geq -1$ ,  $\varphi(s) = 0$  for  $s \leq -2$ , and for  $n \in \mathbb{N}$ ,  $u \in E$  let  $\varphi_n(u) = \varphi((I_\mu(u) - c_{\mu_n})/(\mu_n - \mu))$ . Choose  $\gamma_n \in P$  satisfying (5) and define

$$\tilde{\gamma}_n(t) = \gamma_n(t) - \sqrt{\mu_n - \mu} \cdot \varphi_n(\gamma_n(t)) \frac{I'_\mu(\gamma_n(t))}{\|I'_\mu(\gamma_n(t))\|}.$$

Note that (6) holds true for any  $u = \gamma_n(t_n)$  with  $I_{\mu_n}(u) \geq c_{\mu_n} - 2(\mu_n - \mu)$ , and hence (7) is valid for such  $u$ , if  $n \geq n_0$ . Moreover, (6) also implies  $|I_{\mu}(u) - c_{\mu}| < 2\delta$  and thus  $\|I'_{\mu}(u)\| \geq 2\delta$ .

By (7) and Lemma 3.4 ii), for such  $u$  and sufficiently large  $n \geq n_0$  we also obtain

$$\begin{aligned} \langle I'_{\mu_n}(u), I'_{\mu}(u) \rangle &= \|I'_{\mu}(u)\|^2 - \langle I'_{\mu}(u) - I'_{\mu_n}(u), I'_{\mu}(u) \rangle \geq \\ &\geq \frac{1}{2} \|I'_{\mu}(u)\|^2 - \frac{1}{2} \|I'_{\mu}(u) - I'_{\mu_n}(u)\|^2 \geq \frac{1}{2} \|I'_{\mu}(u)\|^2 - C|\mu - \mu_n|^2 \geq \frac{1}{4} \|I'_{\mu}(u)\|^2 \geq \delta^2. \end{aligned}$$

Thus, by Lemma 3.4 i), for such  $u$ , letting  $\tilde{u} = \tilde{\gamma}_n(t_n)$ ,

$$\begin{aligned} I_{\mu_n}(\tilde{u}) &\leq I_{\mu_n}(u) - \frac{1}{4} \sqrt{\mu_n - \mu} \cdot \varphi_n(u) \|I'_{\mu}(u)\| + \frac{1}{2} |\mu_n - \mu| \varphi_n^2(u) \leq \\ &\leq I_{\mu_n}(u) - \frac{\delta}{4} \sqrt{\mu_n - \mu} \cdot \varphi_n(u) \leq I_{\mu_n}(u) \end{aligned}$$

for  $n \geq n_0$ , and we can estimate

$$\begin{aligned} c_{\mu_n} &\leq \max_{0 \leq t \leq 1} I_{\mu_n}(\tilde{\gamma}_n(t)) = \max_{\{t; I_{\mu_n}(\gamma_n(t)) \geq c_{\mu_n} - (\mu_n - \mu)\}} I_{\mu_n}(\tilde{\gamma}_n(t)) \leq \\ &\leq \max_{0 \leq t \leq 1} I_{\mu_n}(\gamma_n(t)) - \frac{\delta}{4} \sqrt{\mu_n - \mu} \leq \max_{0 \leq t \leq 1} I_{\mu}(\gamma_n(t)) - \frac{\delta}{4} \sqrt{\mu_n - \mu} \leq \\ &\leq c_{\mu} + (\mu_n - \mu) - \frac{\delta}{4} \sqrt{\mu_n - \mu} \leq c_{\mu_n} + \alpha(\mu_n - \mu) - \frac{\delta}{4} \sqrt{\mu_n - \mu} < c_{\mu_n} \end{aligned}$$

for  $n \geq n_0$ , giving the desired contradiction. ■

PROOF OF LEMMA 3.3 (completed). – Let  $(u_n)$  be a sequence as determined in Lemma 3.6. We may assume that  $u_n \rightarrow u$  weakly in  $E$  as  $n \rightarrow \infty$ , and  $e^{u_n} \rightarrow e^u$  in  $L^2$ . Thus,

$$o(1) = \langle I'_{\mu}(u_n), u_n - u \rangle = \|u_n - u\|^2 - o(1),$$

where  $o(1) \rightarrow 0$  as  $\varepsilon \rightarrow 0$ . The claim follows. ■

By Lemma 3.3 problem  $(3)_{\lambda}$  admits a non-trivial solution for almost every  $\lambda \in ]8\pi, 4\pi^2[$ . We now show that this is in fact true for *all*  $\lambda$  in this range.

#### 4. – Compactness.

Theorem 2.1 will be a consequence of Lemma 3.3 and the following compactness result.

LEMMA 4.1. – Let  $\lambda_n \rightarrow \lambda$  and let  $u_n \in E$  be a solution for  $(\mathfrak{S})_{\lambda_n}$ . If  $\lambda \neq 8\pi m$ ,  $m \in \mathbb{N}$ , then  $u_n$  admits a subsequence which converges smoothly to a solution of  $(\mathfrak{S})_\lambda$ .

PROOF. – Our proof relies on a result of Brezis-Merle [1] and its completion given by Li-Shafrir [8]. ■

THEOREM 4.2 (Brezis-Merle). – Let  $D$  be a bounded domain in  $\mathbb{R}^2$  and  $\{w_n\}$  be a sequence satisfying:

$$-\Delta w_n = V_n(x) e^{w_n} \quad \text{on } D$$

with  $0 \leq V_n(x) \leq b_1$  on  $D$ . Also suppose that  $\int_D e^{w_n} \leq b_2$ . Then  $\{w_n\}$  admits a subsequence  $\{w_{n_k}\}$  satisfying one of the following:

- i)  $\{w_{n_k}\}$  is uniformly locally bounded in  $D$ ;
- ii) for any compact set  $K \subset D$ , there holds

$$\sup_K w_{n_k} \rightarrow -\infty \quad \text{as } k \rightarrow +\infty;$$

iii) there exists  $S = \{a_1, \dots, a_p\} \subset D$  (blow up set) and a sequence  $\{x_{n_k}^i\} \subset D$  such that, as  $k \rightarrow \infty$ ,  $x_{n_k}^i \rightarrow a_i$ ,  $w_{n_k}(x_{n_k}^i) \rightarrow \infty$ ,  $i = 1, \dots, p$ .

Moreover, for any compact set  $K \subset D \setminus S$  we have,  $\sup_K w_{n_k} \rightarrow -\infty$  as  $k \rightarrow \infty$ . (Li-Shafrir): In addition, if  $V_n \rightarrow V$  in  $C^0(\bar{\Omega})$ , then  $K$

$$V_{n_k} e^{w_{n_k}} \rightarrow \sum_{i=1}^p 8\pi m_i \delta_{x=a_i}$$

in the sense of measures, with  $m_i \in \mathbb{N}$  and  $\delta_{x=a_i}$  the Dirac distribution supported in  $\{a_i\}$ ,  $i = 1, \dots, p$ .

Theorem 4.2 implies Lemma 4.1, as follows.

PROOF OF LEMMA 4.1. – After translation we may assume that

$$u_n(0) = \sup_\Omega u_n.$$

Let

$$w_n(x) = u_n(x) - \ln \left( \int_\Omega e^{u_n} dx \right) - \frac{\lambda_n}{4} |x|^2,$$

satisfying

$$-\Delta w_n = -\Delta u_n + \lambda_n = \lambda_n \frac{e^{u_n}}{\int_\Omega e^{u_n} dx} = \lambda_n e^{(\lambda_n/4)|x|^2} e^{w_n}$$

with

$$\int_{\Omega} e^{w_n} dx \leq 1 .$$

Thus, the hypotheses of Theorem 4.2 are satisfied for  $w_n$  with  $V_n = \lambda_n e^{(\lambda_n/4)|x|^2} \leq \lambda_n e^{\lambda_n}$ ,  $b_2 = 1$ .

By Theorem 4.2, passing to a sub-sequence if necessary,  $(w_n)$  satisfies (i), (ii) or (iii). Suppose alternative (iii) of Theorem 4.2 holds. Since  $V_n \rightarrow \lambda e^{(\lambda/4)|x|^2}$ , this implies

$$\lambda_n = \int_{\Omega} V_n e^{w_n} \rightarrow \bar{\lambda} \in 8\pi\mathbb{N} \quad \text{as } n \rightarrow \infty .$$

But  $\lambda_n \rightarrow \lambda \in ]8\pi, 4\pi^2[$ , showing that (iii) cannot occur.

Consequently, there exists  $C$  such that

$$C \geq \sup_{B_{1/2}(0)} w_n \geq \sup_{B_{1/2}(0)} u_n - \ln \left( \int_{\Omega} e^{u_n} dx \right) - \frac{\lambda_n}{8} ,$$

and we conclude that

$$u_n(0) - \ln \left( \int_{\Omega} e^{u_n} dx \right) = \sup_{\Omega} u_n - \ln \left( \int_{\Omega} e^{u_n} dx \right) \leq C .$$

Hence  $\Delta u_n \in L^{\infty}(\Omega)$  with

$$\sup_{\Omega} |\Delta u_n| \leq \lambda_n \sup_{\Omega} \left( \frac{e^{u_n}}{\int_{\Omega} e^{u_n} dx} + 1 \right) \leq \lambda_n (e^C + 1) .$$

Consequently, there exists a constant  $C_2$  such that

$$\|u_n\|_{C^{1,\alpha}(\bar{\Omega})} \leq C_2 \quad \text{for all } n \in \mathbb{N}$$

and any fixed  $\alpha \in ]0, 1[$ . Therefore,  $(u_n)$  admits a subsequence which converges in  $C^1(\bar{\Omega})$ —hence smoothly—to a solution of  $(3)_{\lambda}$ . ■

PROOF OF THEOREM 2.1. – Fix  $\lambda \in ]8\pi, 4\pi^2[$ . By Lemmas 3.3 and 4.1 there exists a sequence  $(\lambda_n)$  of numbers  $\lambda_n \leq \lambda$  and corresponding solutions  $u_n$  of  $(3)_{\lambda_n}$ ,  $u$  of  $(3)_{\lambda}$  such that  $I_{\lambda_n}(u_n) = c_{\lambda_n}$ ,  $\lambda_n \rightarrow \lambda$ , and  $u_n \rightarrow u$  smoothly as  $n \rightarrow \infty$ . Since  $c_{\lambda} \leq c_{\lambda_n}$  for all  $n$ , we conclude that

$$I_{\lambda}(u) = \lim_{n \rightarrow \infty} I_{\lambda_n}(u_n) \geq c_{\lambda} > 0 ,$$

showing that  $u \neq 0$ . ■

**5. – Nonexistence for small  $\lambda > 0$ .**

Although we cannot say at this stage whether the «branch» of non-trivial solutions to  $(3)_\lambda$  constructed in Theorem 2.1 extends to  $\lambda \leq 8\pi$ , we can exclude the existence of non-trivial solutions to  $(3)_\lambda$  for small  $\lambda > 0$ .

LEMMA 5.1. – *There exists a constant  $C$  such that for any solution  $u$  to  $(3)_\lambda$  with  $0 \leq \lambda < 4\pi$  there holds*

$$\sup_{\Omega} |u| + \|u\|^2 \leq C \left( \lambda + \left( \frac{\lambda}{4\pi - \lambda} \right)^2 \right).$$

PROOF. – Let  $G$  be the Green’s function to  $-\Delta$  on  $\Omega$ , satisfying  $\int_{\Omega} G(x, y) dy = 0$  for all  $x$ . We have

$$G(x, y) = \frac{1}{2\pi} \ln \frac{1}{|x - y|} + \gamma(x, y),$$

where  $\gamma$ , the regular part of  $G$ , is smooth on  $\bar{\Omega} \times \bar{\Omega}$ .

Then for any  $y \in \Omega$  we find

$$\begin{aligned} (8) \quad u(y) &= - \int_{\Omega} \Delta u G(x, y) dx = \lambda \frac{\int_{\Omega} e^u G(x, y) dx}{\int_{\Omega} e^u dx} \leq \\ &\leq \frac{\lambda}{2\pi} \frac{\int_{\Omega} \ln(1/|x - y|) e^u dx}{\int_{\Omega} e^u dx} + \lambda \|\gamma\|_{L^\infty}. \end{aligned}$$

Using the inequality

$$ab \leq e^a + b(\ln b - 1) \quad \text{for } b > 0, \quad a \in \mathbb{R},$$

which follows from the equation

$$\sup_a \{ab - e^a\} = b(\ln b - 1),$$

and letting  $a = \alpha \ln \left( \frac{1}{|x - y|} \right) = \ln \left( \frac{1}{|x - y|^\alpha} \right)$ ,  $b = \frac{e^u}{\alpha}$  for  $1 \leq \alpha < 2$ , the first term may be estimated

$$\begin{aligned} (9) \quad \frac{\int_{\Omega} \ln(1/|x - y|) e^u dx}{\int_{\Omega} e^u dx} &\leq \frac{\int_{\Omega} (1/|x - y|^\alpha) dx}{\int_{\Omega} e^u dx} + \frac{\int_{\Omega} e^u u dx}{\alpha \int_{\Omega} e^u dx} + C \leq \\ &\leq \frac{C}{2 - \alpha} + \frac{\int_{\Omega} e^u u dx}{\alpha \int_{\Omega} e^u dx}, \end{aligned}$$

for any  $1 \leq \alpha < 2$ , where we also used that  $\int_{\Omega} e^u dx \geq 1$ . Now observe that

$$\|u\|^2 = \lambda \frac{\int_{\Omega} e^u u dx}{\int_{\Omega} e^u dx}.$$

Together with the above estimate this implies that

$$\|u\|^2 \leq \lambda \sup_{\Omega} u \leq \frac{C\lambda^2}{2-\alpha} + \frac{\lambda^2}{2\pi} \frac{\int_{\Omega} e^u u dx}{\alpha \int_{\Omega} e^u dx} = \frac{C\lambda^2}{2-\alpha} + \frac{\lambda}{2\pi\alpha} \|u\|^2.$$

Thus for  $\lambda < 4\pi$  we obtain the estimate

$$\|u\|^2 \leq \inf_{1 \leq \alpha < 2} \frac{C\lambda^2}{(2-\alpha)(2\pi\alpha - \lambda)} \leq C \left( \frac{\lambda}{4\pi - \lambda} \right)^2$$

with a uniform constant  $C$ .

From (8) and (9) with  $\alpha = 1$  we then also derive that

$$\sup_{\Omega} |u| \leq C\lambda + \frac{1}{2\pi} \|u\|^2 \leq C\lambda + C \left( \frac{\lambda}{4\pi - \lambda} \right)^2,$$

as claimed. ■

**THEOREM 5.10.** – *There exists  $A > 0$  such that for  $0 \leq \lambda < A$  any solution  $u \in E$  of  $(3)_{\lambda}$  vanishes identically.*

**PROOF.** – By Lemma 5.1 for any solution  $u$  of  $(3)_{\lambda}$  for  $0 \leq \lambda \leq A < 4\pi$  we can bound

$$\sup_{\Omega} |u| \leq C\lambda$$

with a constant  $C = C(A)$ . Thus, we also have

$$|e^u - 1| \leq e^{C\lambda} u.$$

Since  $\int_{\Omega} u dx = 0$ ,  $\int_{\Omega} e^u dx \geq 1$ , it follows that

$$\|u\|^2 = \lambda \frac{\int_{\Omega} e^u u dx}{\int_{\Omega} e^u dx} = \lambda \frac{\int_{\Omega} (e^u - 1) u dx}{\int_{\Omega} e^u dx} \leq \lambda e^{C\lambda} \int_{\Omega} u^2 dx \leq \frac{\lambda e^{C\lambda}}{4\pi^2} \|u\|^2,$$

and the claim follows. ■

## REFERENCES

- [1] H. BREZIS - F. MERLE, *Uniform estimates and blow-up behavior for solutions of  $-\Delta u = V(x)e^u$  in two dimensions*, Comm. P.D.E., **16** (1991), 1223-1253.
- [2] E. CAGLIOTI - P. L. LIONS - C. MARCHIORO - M. PULVIRENTI, *A special class of stationary flows for two-dimensional Euler equations, a statistical mechanics description*, Comm. Math. Phys., **143** (1992), 501-525.
- [3] E. CAGLIOTI - P. L. LIONS - C. MARCHIORO - M. PULVIRENTI, *A special class of stationary flows for two-dimensional Euler equations, a statistical mechanics description, part II*, Comm. Math. Phys., **174** (1995), 229-260.
- [4] G. DUNNE, *Self-dual Chern Simons Theories*, Lecture Notes in Physics, New Series M, **36**, Springer, New York (1996).
- [5] J. HONG - Y. KIM - P. Y. PAC, *Multivortex solutions of the Abelian Chern Simons theory*, Phys. Rev. Lett., **64** (1990), 2230-2233.
- [6] R. JACKIW - E. J. WEINBERG, *Selfdual Chern Simons vortices*, Phys. Rev. Lett., **64** (1990), 2234-2237.
- [7] M. K. H. KIESSLING, *Statistical Mechanics of classical particles with logarithmic interaction*, Comm. Pure Appl. Math., **46** (1993), 27-56.
- [8] Y. LI - I. SHAFRIR, *Blow-up analysis for solutions of  $-\Delta u = Ve^u$  in dimension two*, Ind. Univ. Math. J., **43** (1994), 1255-1270.
- [9] J. MOSER, *A sharp form of an inequality by N. Trudinger*, Ind. Univ. Math. J., **20** (1971), 1077-1091.
- [10] T. RICCIARDI - G. TARANTELO, in preparation.
- [11] M. STRUWE, *The existence of surfaces of constant mean curvature with free boundaries*, Acta Math., **160** (1988), 19-64.
- [12] M. STRUWE, *Critical points of embeddings of  $H^{1,2}$  into Orlicz spaces*, Ann. Inst. H. Poincaré, Analyse Nonlin., **5** (1988), 425-464.
- [13] T. SUZUKI, *Global analysis for two dimensional elliptic eigenvalue problems with exponential nonlinearities*, Ann. Inst. H. Poincaré, Analyse Nonlin., **9** (1992), 367-398.
- [14] G. TARANTELO, *Multiple condensate solutions for the Chern-Simons-Higgs theory*, J. Math. Phys., **37** (8) (1996), 3769-3796.
- [15] C. H. TAUBES, *Arbitrary  $N$ -vortex solutions to the first order Ginzburg-Landau equation*, Comm. Math. Phys., **72** (1980), 277-292.

M. Struwe: D-Math, ETH-Zentrum, CH-8092 Zürich, Switzerland  
 e-mail: struwe@math.ethz.ch

G. Tarantello: Università di Roma «Tor Vergata», Dipartimento di Matematica  
 Via della Ricerca Scientifica, 00133 Roma, Italy  
 e-mail: tarantello@mat.utovrm.it