

Universität Karlsruhe

Institut für Analysis

Prof. Dr. Michael Plum

Dipl.-phys. Vu Hoang

Boundary and Eigenvalue Problems 3rd Exercise Sheet

- 7) a) Let $L[u] = (-\Delta)^k u = \underbrace{(-\Delta) \circ \dots \circ (-\Delta)}_{k\text{-times}} u$. Show that L is uniformly strongly elliptic.

- b) Show that

$$L[u] = \operatorname{div} \left(\frac{\nabla u}{\sqrt{1 + |\nabla u|^2}} \right)$$

is a quasilinear elliptic differential operator.

- 8) Let $\Omega = (0, 1) \times (0, 1) \subset \mathbb{R}^2$ and consider the boundary value problem

$$(1) \quad \begin{cases} u_{xy} = 0 & \text{in } \Omega \\ u = u_0 & \text{on } \partial\Omega \end{cases}$$

for $u \in C^2(\Omega) \cap C(\bar{\Omega})$ and given $u_0 \in C(\partial\Omega)$. Prove that the relation

$$u(0, 0) + u(1, 1) = u(1, 0) + u(0, 1)$$

holds for any solution of (1) and conclude that there exists a $u_0 \in C(\partial\Omega)$ such that (1) has no solution. Is the operator $L[u] = u_{xy}$ elliptic?

- 9) Let $\Omega \subset \mathbb{R}^n$ be an open, bounded set with Lipschitz-boundary. Verify for $u, v \in C^1(\bar{\Omega}) \cap C^2(\Omega)$ the (generalized) Green's identities

$$\text{a) } \int_{\Omega} L[u]v dx = - \int_{\partial\Omega} \frac{\partial u}{\partial \mu} v d\sigma + \int_{\Omega} (\nabla u)^T A \nabla v dx$$

$$\text{b) } \int_{\Omega} (L[u]v - uL[v]) dx = \int_{\partial\Omega} \left(\frac{\partial v}{\partial \mu} u - v \frac{\partial u}{\partial \mu} \right) dx$$

where $L[u] = - \sum_{i,j=1}^m \frac{\partial}{\partial x_i} \left(a_{ij} \frac{\partial u}{\partial x_j} \right)$ ($a_{ij} \in C^1(\bar{\Omega})$, $A(x) = (a_{ij}(x))$ symmetric matrix).