

## Classical Methods for Partial Differential Equations

### Exercise sheet 8

#### Exercise 26

Let  $\Omega \subseteq \mathbb{R}^n$  be a domain,  $n \geq 2$ ,  $x_0 \in \Omega$  and  $u: \Omega \setminus \{x_0\} \rightarrow \mathbb{R}$  be a harmonic function. We say that the point  $x_0$  is a *removable singularity of the function*  $u$ , if there exists a harmonic function  $\tilde{u}: \Omega \rightarrow \mathbb{R}$  such that  $\tilde{u}(x) = u(x)$  ( $x \in \Omega \setminus \{x_0\}$ ). Let  $S$  denote the usual singularity function defined in the lecture. Show that

1. If  $\lim_{x \rightarrow x_0} \frac{u(x)}{S(x_0, x)} = 0$ , then  $x_0$  is a removable singularity of  $u$ .
2. If  $\lim_{x \rightarrow x_0} \frac{u(x)}{S(x_0, x)} = c$ , for some  $c \in \mathbb{R}$ , then there exists a harmonic function  $v: \Omega \rightarrow \mathbb{R}$  such that  $u(x) = cS(x_0, x) + v(x)$  ( $x \in \Omega \setminus \{x_0\}$ ).

#### Exercise 27

Let  $x_0 \in \mathbb{R}^n$ ,  $R > 0$  and  $\Omega \subseteq \mathbb{R}^n$  be a domain, such that  $\Omega \subseteq B_R(x_0)$ . Moreover, let  $q \in \mathcal{C}(\overline{\Omega})$ ,  $\varphi \in \mathcal{C}(\partial\Omega)$  and

$$\overline{K} = \max_{x \in \overline{\Omega}} \{q(x), 0\}, \quad \underline{K} = \min_{x \in \overline{\Omega}} \{q(x), 0\}, \quad \overline{M} = \max_{x \in \partial\Omega} \varphi(x), \quad \underline{M} = \min_{x \in \partial\Omega} \varphi(x).$$

Show that if  $u \in \mathcal{C}^2(\Omega) \cap \mathcal{C}(\overline{\Omega})$  is a solution of the following boundary value problem

$$\begin{cases} -\Delta u = q & \text{in } \Omega, \\ u = \varphi & \text{on } \partial\Omega, \end{cases}$$

then the following inequality is true

$$\underline{M} + \frac{\underline{K}}{2n}(R^2 - |x - x_0|^2) \leq u(x) \leq \overline{M} + \frac{\overline{K}}{2n}(R^2 - |x - x_0|^2).$$

#### Exercise 28

Prove the Harnack inequality: Let  $\Omega = \{x \in \mathbb{R}^n: |x| < R\}$  and  $u \in \mathcal{C}^2(\Omega) \cap \mathcal{C}(\overline{\Omega})$ . If  $u$  is harmonic in  $\Omega$  and  $u(x) \geq 0$  ( $x \in \Omega$ ), then

$$\frac{1 - \frac{|x|}{R}}{\left(1 + \frac{|x|}{R}\right)^{n-1}} u(0) \leq u(x) \leq \frac{1 + \frac{|x|}{R}}{\left(1 - \frac{|x|}{R}\right)^{n-1}} u(0) \quad (x \in \Omega).$$

### Exercise 29

Let  $R > 0$ ,  $\Omega = \mathbb{R}^n \setminus \overline{B_R(0)}$  and  $u \in \mathcal{C}^2(\Omega) \cap \mathcal{C}(\overline{\Omega})$  be a harmonic function such that  $\lim_{|x| \rightarrow \infty} u(x) = 0$ .

1. Show that  $\sup_{x \in \Omega} |u(x)| = \max_{x \in \partial\Omega} |u(x)|$ .
2. Find an example showing that the above statement is false, if the assumption  $\lim_{|x| \rightarrow \infty} u(x) = 0$  is not satisfied.

Above exercises will be discussed on 16.12.2015.