

## Classical Methods for Partial Differential Equations

### Exercise sheet 9

#### Exercise 30

Let  $\Omega \subseteq \mathbb{R}^n$  be a domain and let  $N \subseteq \mathbb{R}^n$  be an open neighbourhood of a point  $\xi \in \partial\Omega$ . A function  $w: \overline{\Omega} \cap N \rightarrow \mathbb{R}$  is called a *local barrier function* at the point  $\xi$  if the following conditions are satisfied

1.  $w \in \mathcal{C}(\overline{\Omega} \cap N)$  is subharmonic in  $\Omega \cap N$ ,
2.  $w(x) < 0$  ( $x \in (\overline{\Omega} \cap N) \setminus \{\xi\}$ ),
3.  $w(\xi) = 0$ .

Show what if for a point  $\xi \in \partial\Omega$  there exists a local barrier function, then there exists a barrier function at this point.

*Hint:* Let  $B_\rho(\xi)$  be a ball around  $\xi$  with sufficiently small radius  $\rho > 0$  and  $m \in \mathbb{R}$  be chosen in a suitable way. Consider a function

$$\bar{w}(x) = \begin{cases} \max\{m, w(x)\} & (x \in \overline{\Omega} \cap B_\rho(\xi)), \\ m & (x \in \overline{\Omega} \setminus B_\rho(\xi)). \end{cases}$$

#### Exercise 31

Let  $\Omega \subseteq \mathbb{R}^n$  be a domain satisfying the outer ball condition. Show that for each point of the boundary  $\partial\Omega$  a barrier function exists.

*Hint:* Singularity function.

#### Exercise 32

A set  $K \subseteq \mathbb{R}^2$  is called a *cone with an opening angle*  $\alpha \in (0, 2\pi)$ , if  $K$  can be transformed by rigid motion (rotations and translations) to a set

$$K_0 = \{(r \cos \varphi, r \sin \varphi) \in \mathbb{R}^2 : 0 < r < \infty, 0 < \varphi < \alpha\}.$$

Let  $\Omega \subseteq \mathbb{R}^2$  be a domain and let  $\xi \in \partial\Omega$ . Assume that there exists a cone  $K$  such that  $K \subseteq \mathbb{R}^2 \setminus \Omega$  and  $\overline{K} \cap \overline{\Omega} = \{\xi\}$ . Show that there exists a local barrier function at the point  $\xi$ .

*Hint:* Consider a function  $w(r, \varphi) = r^{\frac{\pi}{2\pi-\alpha}} \sin\left(\frac{\pi}{2\pi-\alpha}\varphi\right)$ .

### Exercise 33

Let  $\Omega \subseteq \mathbb{R}^n$  be a domain,  $M > 0$  and  $(u_j)_{j \in \mathbb{N}} \subseteq \mathcal{C}(\overline{\Omega})$  be a sequence of functions which are harmonic in  $\Omega$  and satisfy

$$|u_j(x)| \leq M \quad (x \in \overline{\Omega}, j \in \mathbb{N}).$$

Show that there exists a subsequence  $(u_{j_k})_{k \in \mathbb{N}}$  and a function  $u \in \mathcal{C}(\Omega)$ , such that for all open and bounded sets  $U \subseteq \Omega$  such that  $\overline{U} \subseteq \Omega$

$$u_{j_k} \xrightarrow[k \rightarrow \infty]{} u \text{ uniformly in } U.$$

*Hint:* Apply the gradient estimate in theorem 3.13 from the lecture and the Arzelà–Ascoli theorem.

Above exercises will be discussed on 21.12.2015.