



On a Generalized Punctured Neighborhood Theorem in $\mathscr{L}(X)$

Christoph Schmoeger

Proceedings of the American Mathematical Society, Vol. 123, No. 4. (Apr., 1995), pp. 1237-1240.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9939%28199504%29123%3A4%3C1237%3AOAGPNT%3E2.0.CO%3B2-6>

Proceedings of the American Mathematical Society is currently published by American Mathematical Society.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ams.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is an independent not-for-profit organization dedicated to creating and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact support@jstor.org.

ON A GENERALIZED PUNCTURED NEIGHBORHOOD THEOREM IN $\mathcal{L}(X)$

CHRISTOPH SCHMOEGER

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. Suppose that T is a bounded linear operator on a complex Banach space X . If $T^2(X)$ is closed, $T(X) \cap N(T)$ is finite dimensional, and S is a bounded linear operator on X such that S is invertible, commutes with T , and has sufficiently small norm, then $T - S$ is upper semi-Fredholm.

Throughout this paper X will denote a complex Banach space. We write $\mathcal{L}(X)$ for the set of all bounded linear operators on X . For $T \in \mathcal{L}(X)$, we denote by $N(T)$ the kernel and by $T(X)$ the range of T . The operator T is called *upper semi-Fredholm* if $T(X)$ is closed and $\dim N(T) < \infty$. We write $\sigma(T)$ for the spectrum of T . It is well known that the resolvent $R_\lambda(T) = (\lambda I - T)^{-1}$ is a holomorphic function of λ for points λ in the resolvent set $\mathbb{C} \setminus \sigma(T)$.

The aim of this paper is the following generalization of the “punctured neighborhood theorem” for upper semi-Fredholm operators:

Theorem 1. *Suppose that $T \in \mathcal{L}(X)$, T^2 has closed range, and $T(X) \cap N(T)$ is finite dimensional. Then:*

(a) *$T - S$ is upper semi-Fredholm whenever $S \in \mathcal{L}(X)$ is invertible, $TS = ST$, and $\|S\|$ is sufficiently small. Furthermore, we have*

$$\dim N(T - S) = \dim \left(N(T) \cap \bigcap_{n=1}^{\infty} T^n(X) \right).$$

(b) *If 0 is a boundary point of $\sigma(T)$, then 0 is a pole of the resolvent of T .*

For the proof of Theorem 1 we need some additional notation and a preliminary lemma.

Let $T \in \mathcal{L}(X)$. We write $\alpha(T)$ and $\beta(T)$ for $\dim N(T)$ and $\text{codim } T(X)$, respectively. The operator T is called *lower semi-Fredholm* if $\beta(T)$ is finite (in this case T has closed range, by [4, Satz 55.4]). T is called *semi-Fredholm* if T is upper or lower semi-Fredholm. T is *Fredholm* if both $\alpha(T)$ and $\beta(T)$ are finite. The *index* of a semi-Fredholm operator T is defined by

$$\text{ind}(T) = \alpha(T) - \beta(T).$$

Received by the editors November 9, 1992 and, in revised form, August 2, 1993.

1991 *Mathematics Subject Classification.* Primary 47A10, 47A53, 47A55.

Key words and phrases. Semi-Fredholm operator, spectrum, essential spectrum.

For $T \in \mathcal{L}(X)$ we define the number $k_n(T)$ by

$$k_n(T) = \dim((N(T) \cap T^n(X))/(N(T) \cap T^{n+1}(X))) \quad (n \geq 0).$$

We say that T has *uniform descent* for $n \geq d$ if $k_n(T) = 0$ for $n \geq d$. This notion is due to Grabiner (see [3, Definition 1.3 and Lemma 2.3]). If T has uniform descent for $n \geq d$ and if $T^n(X)$ is closed in the operator range topology of $T^d(X)$ for $n \geq d$, then we say that T has *topological uniform descent* for $n \geq d$ (see [3, Definition 2.5]). For a discussion of operator ranges and their topologies, the reader is referred to [1] or [2].

Lemma. *Suppose that $T \in \mathcal{L}(X)$, T^2 has closed range, and $\dim T(X) \cap N(T) < \infty$. Then:*

- (a) *There exists an integer $d \geq 0$ such that T has uniform descent for $n \geq d$.*
- (b) *T has topological uniform descent for $n \geq d$ and $\bigcap_{n=1}^{\infty} T^n(X)$ is closed.*
- (c) *$T(\bigcap_{n=1}^{\infty} T^n(X)) = \bigcap_{n=1}^{\infty} T^n(X)$.*
- (d) *If $S \in \mathcal{L}(X)$ is invertible, $TS = ST$, and $\|S\|$ is sufficiently small, then $T - S$ has closed range.*

Proof. (a) Since

$$N(T) \cap T^{n+1}(X) \subseteq N(T) \cap T^n(X) \subseteq N(T) \cap T(X) \quad \text{for } n \geq 0$$

and $\dim(N(T) \cap T(X)) < \infty$, the result follows.

(b) Invoke [3, Lemma 2.4]. The hypotheses of this lemma are satisfied because of [3, Lemma 2.3].

(c) Use (b) and [3, Theorem 3.4(a)].

(d) Put $V = T - S$. Since T has topological uniform descent for $n \geq d$, V has closed range, by [3, Theorem 4.7(a)]. \square

Proof of Theorem 1. (a) Put $X_0 = \bigcap_{n=1}^{\infty} T^n(X)$, and denote the restriction of T to X_0 by T_0 . Clearly,

$$\alpha(T_0) = \dim \left(N(T) \cap \bigcap_{n=1}^{\infty} T^n(X) \right) \leq \dim(N(T) \cap T(X)) < \infty.$$

Part (c) of the above lemma shows that $T_0(X_0) = X_0$; thus, $\beta(T_0) = 0$. It follows that T_0 is Fredholm with

$$\text{ind}(T_0) = \alpha(T_0) - \beta(T_0) = \alpha(T_0).$$

[4, Satz 82.4] shows that there exists $\varepsilon > 0$ such that

$$T_0 - R \text{ is Fredholm}$$

and

$$\text{ind}(T_0 - R) = \text{ind}(T_0) \quad \text{for } R \in \mathcal{L}(X_0) \text{ with } \|R\| < \varepsilon.$$

Furthermore, again by [4, Satz 82.4],

$$\alpha(T_0 - R) \leq \alpha(T_0), \quad \beta(T_0 - R) \leq \beta(T_0) = 0$$

for $R \in \mathcal{L}(X_0)$ with $\|R\| < \varepsilon$. This gives $\beta(T_0 - R) = 0$ and

$$(*) \quad \alpha(T_0 - R) = \text{ind}(T_0 - R) = \text{ind}(T_0) = \alpha(T_0)$$

for each $R \in \mathcal{L}(X_0)$ such that $\|R\| < \varepsilon$.

Now suppose that $S \in \mathcal{L}(X)$ is invertible, $TS = ST$, and $\|S\| < \varepsilon$. Since S commutes with T , we have $S(X_0) \subseteq X_0$. Put $S_0 = S|_{X_0}$; then $S_0 \in \mathcal{L}(X_0)$ and $\|S_0\| < \varepsilon$.

Next we show that $N(T - S) = N(T_0 - S_0)$. The inclusion $N(T_0 - S_0) \subseteq N(T - S)$ is clear. Let $x \in N(T - S)$; thus, $Tx = Sx$. Since $TS = ST$, we have $T^n x = S^n x$ for each $n \in \mathbb{N}$; therefore, $x = S^{-n} T^n x = T^n (S^{-n} x) \in T^n(X)$ for each $n \in \mathbb{N}$ and, hence, $x \in X_0$. This proves $N(T - S) = N(T_0 - S_0)$. Therefore, by (*),

$$\alpha(T - S) = \alpha(T_0 - S_0) = \alpha(T_0).$$

To complete the proof of (a), we have to show that $T - S$ has closed range for sufficiently small $\|S\|$. But this follows immediately from part (d) of the above lemma.

(b) follows from [3, Corollary 4.9]. \square

Remark. Theorem 1 can be proved under the weaker assumption that $T^{-1}(T^2(X)) = T(X) + N(T)$ is closed. $T(X) + N(T)$ being closed is equivalent to $T^2(X)$ is closed in the operator range topology on $T(X)$ (see the proof of [3, Theorem 3.2]), so our results are still true except that $X_0 = \bigcap_{n=1}^{\infty} T^n(X)$ is only known to be closed in the operator range topology on $T(X)$.

For $T \in \mathcal{L}(X)$ we define two essential spectra:

$$\sigma_e(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not Fredholm}\}$$

and

$$\sigma_w(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not a Fredholm operator with } \text{ind}(T - \lambda I) = 0\}.$$

Theorem 2. *Let $T \in \mathcal{L}(X)$, and suppose that $T^2(X)$ is closed and $\dim N(T) \cap T(X) < \infty$. Then:*

- (a) *If 0 is a boundary point of $\sigma_e(T)$, 0 is isolated in $\sigma_e(T)$.*
- (b) *If 0 is a boundary point of $\sigma_w(T)$, 0 is isolated in $\sigma_w(T)$.*

Proof. Since T has topological uniform descent for $n \geq d$ (Lemma, part (b)), we can define

$$\alpha^*(T) = \lim_{n \rightarrow \infty} \dim N(T^{n+1})/N(T^n)$$

and

$$\beta^*(T) = \lim_{n \rightarrow \infty} \dim T^n(X)/T^{n+1}(X).$$

Theorem 4.7 in [3] then says that if $\lambda \in \mathbb{C} \setminus \{0\}$ is sufficiently small, then $\alpha(T - \lambda I) = \alpha^*(T - \lambda I) = \alpha^*(T)$ and $\beta(T - \lambda I) = \beta^*(T - \lambda I) = \beta^*(T)$. Thus, if any $T - \lambda I$ is semi-Fredholm for small $\lambda \neq 0$, then all are semi-Fredholm with the same index $\alpha^*(T) - \beta^*(T)$. This gives the results. \square

ACKNOWLEDGMENT

The author thanks the referee, whose suggestions led to an improvement of the paper.

REFERENCES

1. P. A. Fillmore and J. P. Williams, *On operator ranges*, Adv. Math. **7** (1971), 253–281.
2. S. Grabiner, *Ranges of products of operators*, Canad. J. Math. **26** (1974), 1430–1441.
3. —, *Uniform ascent and descent of bounded operators*, J. Math. Soc. Japan **34** (1982), 317–337.
4. H. Heuser, *Funktionalanalysis*, 2nd ed., Teubner, Stuttgart, 1986.

MATHEMATISCHES INSTITUTE I, UNIVERSITÄT KARLSRUHE, POSTFACH 6980, D-76128 KARLSRUHE, GERMANY